arXiv:1009.2355v2 [math.RT] 26 Nov 2010

QUIVER VARIETIES AND PATH REALIZATIONS ARISING FROM ADJOINT CRYSTALS OF TYPE $A_n^{(1)}$

SEOK-JIN KANG[1] AND EUIYONG PARK[1,2]

Abstract. Let $B(\Lambda_0)$ be the level 1 highest weight crystal of the quantum affine algebra $U_q(A_n^{(1)})$. We construct an explicit crystal isomorphism between the geometric realization $\mathbb{B}(\Lambda_0)$ of $B(\Lambda_0)$ via quiver varieties and the path realization $\mathcal{P}^{\mathrm{ad}}(\Lambda_0)$ of $B(\Lambda_0)$ arising from the adjoint crystal B^{ad}.

Introduction

The theory of *perfect crystals* developed in [7] has a lot of important and interesting applications to the representation theory of quantum affine algebras and the theory of vertex models in mathematical physics. In particular, the crystal $B(\lambda)$ of an integrable highest weight module over a quantum affine algebra can be realized as the crystal $\mathcal{P}^B(\lambda)$ consisting of λ-paths arising from a given perfect crystal B. In [1], Benkart, Frenkel, Kang and Lee gave a uniform construction of level 1 perfect crystals for all quantum affine algebras. These perfect crystals are called the *adjoint crystals* because, when forgetting 0-arrows, they coincide with the direct sums of the trivial crystals and the crystals of adjoint or little adjoint representations of finite dimensional simple Lie algebras.

On the other hand, for a symmetric Kac-Moody algebra $\mathfrak{g}$, Lusztig gave a geometric construction of $U_q^-(\mathfrak{g})$ in terms of perverse sheaves on quiver varieties and introduced the notion of *canonical basis* which yields natural bases for all integrable highest weight modules as well [13, 14]. In [12], Kashiwara and Saito defined a crystal structure on the set $\mathbb{B}(\infty)$ of irreducible components of Lusztig's quiver varieties and showed that $\mathbb{B}(\infty)$ is isomorphic to the crystal $B(\infty)$ of $U_q^-(\mathfrak{g})$. Moreover, in [16, 17], Nakajima defined a new family of quiver varieties associated with a dominant integral weight λ and gave a geometric realization of the integrable highest weight $\mathfrak{g}$-module $V(\lambda)$. In [18], Saito defined a crystal structure on the set $\mathbb{B}(\lambda)$ of irreducible components of certain Lagrangian subvarieties of Nakajima's quiver varieties, and showed that $\mathbb{B}(\lambda)$ is isomorphic to the crystal $B(\lambda)$ of $V(\lambda)$.

Therefore, for quantum affine algebras, it is natural to investigate the crystal isomorphism between the geometric realization $\mathbb{B}(\lambda)$ and the path realization $\mathcal{P}^B(\lambda)$ for various perfect crystals B. In this paper, we will focus on the level 1 highest weight crystal $B(\Lambda_0)$ of the quantum affine algebra $U_q(A_n^{(1)})$, and construct an explicit crystal isomorphism between $\mathbb{B}(\Lambda_0)$ and $\mathcal{P}^{\mathrm{ad}}(\Lambda_0)$, where $\mathcal{P}^{\mathrm{ad}}(\Lambda_0)$ is

---

2000 *Mathematics Subject Classification.* 17B10, 17B67, 81R10.

Key words and phrases. adjoint crystal, crystal graph, Kac-Moody algebra, path realization, quiver variety.

[1] This research was supported by KRF Grant # 2007-341-C00001.

[2] This research was supported by BK21 Mathematical Sciences Division.

[†] This is accepted for publication by the Transactions of the American Mathematical Society.

the path realization arising from the adjoint crystal B^{ad}. We will also give a geometric interpretation of the *fundamental isomorphism theorem for perfect crystals*: $B(\Lambda_0) \cong B(\Lambda_0) \otimes B^{\mathrm{ad}}$. One of the key ingredients of our construction is the explicit 1-1 correspondence between $\mathbb{B}(\Lambda_0)$ and $\mathcal{Y}(\Lambda_0)$ discovered in [3, 19], where $\mathcal{Y}(\Lambda_0)$ is the crystal consisting of $(n+1)$-reduced Young diagrams. We hope our construction will provide a new insight toward the understanding of the connection between $\mathbb{B}(\Lambda_0)$ and $\mathcal{P}^{\mathrm{ad}}(\Lambda_0)$ for *all* quantum affine algebras.

Acknowledgments. The authors wish to express their sincere gratitude to National Institute for Mathematical Sciences in Daejon, Korea, for their generous support during the special program (Thematic Program TP0902) on representation theory in 2009.

1. THE QUANTUM AFFINE ALGEBRA $U_q(A_n^{(1)})$

Let $I = \mathbb{Z}/(n+1)\mathbb{Z}$ be the index set. The *affine Cartan datum of $A_n^{(1)}$-type* consists of

(i) the affine Cartan matrix

$$A = (a_{ij})_{i,j\in I} = \begin{pmatrix} 2 & -1 & 0 & \cdots & -1 \\ -1 & 2 & -1 & \cdots & 0 \\ \vdots & & \ddots & & \vdots \\ 0 & \cdots & -1 & 2 & -1 \\ -1 & 0 & \cdots & -1 & 2 \end{pmatrix},$$

(ii) dual weight lattice $P^\vee = \bigoplus_{i=0}^n \mathbb{Z}h_i \oplus \mathbb{Z}d$,

(iii) affine weight lattice $P = \bigoplus_{i=0}^n \mathbb{Z}\Lambda_i \oplus \mathbb{Z}\delta \subset \mathfrak{h}^*$, where

$$\mathfrak{h} = \mathbb{C} \otimes P^\vee,\ \Lambda_i(h_j) = \delta_{ij},\ \Lambda_i(d) = 0,\ \delta(h_i) = 0,\ \delta(d) = 1\ (i,j \in I),$$

(iv) the set of simple coroots $\Pi^\vee = \{h_i |\ i \in I\}$,

(v) the set of simple roots $\Pi = \{\alpha_i |\ i \in I\}$ given by

$$\alpha_j(h_i) = a_{ij},\ \alpha_j(d) = \delta_{0,j}\ (i,j \in I).$$

The free abelian group $Q = \bigoplus_{i=0}^n \mathbb{Z}\alpha_i$ is called the *root lattice* and the semigroup $Q^+ = \sum_{i=0}^n \mathbb{Z}_{\geq 0}\alpha_i$ is called the *positive root lattice*. For $\alpha = \sum_{i\in I} k_i\alpha_i \in Q^+$, the number $\mathrm{ht}(\alpha) = \sum_{i\in I} k_i$ is called the *height* of α. For $\lambda, \mu \in \mathfrak{h}^*$, we define $\lambda \geq \mu$ if and only if $\lambda - \mu \in Q^+$. The elements in $P^+ = \{\lambda \in P |\ \lambda(h_i) \geq 0,\ i \in I\}$ are called the *dominant integral weights*. Note that the minimal *imaginary root* is given by $\delta = \alpha_0 + \alpha_1 + \cdots + \alpha_n \in Q^+$. The element $c = h_0 + h_1 + \cdots + h_n \in P^\vee$ is called the *canonical central element*.

Given $n \in \mathbb{Z}$ and any symbol q, we define

$$[n]_q = \frac{q^n - q^{-n}}{q - q^{-1}}$$

and set $[0]_q! = 1$, $[n]_q! = [n]_q[n-1]_q \cdots [1]_q$. For $m \geq n \geq 0$, let

$$\begin{bmatrix} m \\ n \end{bmatrix}_q = \frac{[m]_q!}{[n]_q![m-n]_q!}.$$

Definition 1.1. The *quantum affine algebra* $U_q(\mathfrak{g}) = U_q(A_n^{(1)})$ is an associative algebra over $\mathbb{C}(q)$ with 1 generated by $e_i,\ f_i\ (i \in I)$ and $q^h\ (h \in P^\vee)$ satisfying the defining relations:

(a) $q^0 = 1,\ q^h q^{h'} = q^{h+h'}$ for $h, h' \in P^\vee$,
(b) $q^h e_i q^{-h} = q^{\alpha_i(h)} e_i$ for $h \in P^\vee$,
(c) $q^h f_i q^{-h} = q^{-\alpha_i(h)} f_i$ for $h \in P^\vee$,
(d) $e_i f_j - f_j e_i = \delta_{ij}(q^{h_i} - q^{-h_i})/(q - q^{-1})$ for $i, j \in I$,
(e) $\sum_{k=0}^{1-a_{ij}} (-1)^k \left[{1-a_{ij} \atop k}\right]_q e_i^{1-a_{ij}-k} e_j e_i^k = 0$ for $i \neq j$,
(f) $\sum_{k=0}^{1-a_{ij}} (-1)^k \left[{1-a_{ij} \atop k}\right]_q f_i^{1-a_{ij}-k} f_j f_i^k = 0$ for $i \neq j$.

The definition of *category* O_{int}^q, *Kashiwara operators* and *crystal bases* can be found in [10, 4], etc. It was shown in [10] that every $U_q(\mathfrak{g})$-module in the category O_{int}^q has a crystal basis. The notion of *abstract crystals* was introduced in [11]. For convenience, we recall some of the basic definitions and properties of abstract crystals.

Definition 1.2. An *abstract crystal* associated with the Cartan datum $(A, \Pi, \Pi^\vee, P, P^\vee)$ is a set B together with the maps $\mathrm{wt} : B \to P,\ \tilde{e}_i, \tilde{f}_i : B \to B \sqcup \{0\}$, and $\varepsilon_i, \varphi_i : B \to \mathbb{Z} \cup \{-\infty\}$ $(i \in I)$ satisfying the following properties:

(a) $\varphi_i(b) = \varepsilon_i(b) + \langle h_i, \mathrm{wt}(b) \rangle$ for all $i \in I$,
(b) $\mathrm{wt}(\tilde{e}_i b) = \mathrm{wt}(b) + \alpha_i$ if $\tilde{e}_i b \in B$,
(c) $\mathrm{wt}(\tilde{f}_i b) = \mathrm{wt}(b) - \alpha_i$ if $\tilde{f}_i b \in B$,
(d) $\varepsilon_i(\tilde{e}_i b) = \varepsilon_i(b) - 1,\ \varphi_i(\tilde{e}_i b) = \varphi_i(b) + 1$ if $\tilde{e}_i b \in B$,
(e) $\varepsilon_i(\tilde{f}_i b) = \varepsilon_i(b) + 1,\ \varphi_i(\tilde{f}_i b) = \varphi_i(b) - 1$ if $\tilde{f}_i b \in B$,
(f) $\tilde{f}_i b = b'$ if and only if $b = \tilde{e}_i b'$ for $b, b' \in B,\ i \in I$,
(g) if $\varphi_i(b) = -\infty$ for $b \in B$, then $\tilde{e}_i b = \tilde{f}_i b = 0$.

We often say that B is a $U_q(\mathfrak{g})$-*crystal*. We denote $B_\lambda = \{b \in B \mid \mathrm{wt}(b) = \lambda\}$ so that $B = \bigsqcup_{\lambda \in P} B_\lambda$.

Example 1.3.

(1) Let (L, B) be a crystal basis of $M \in O_{\mathrm{int}}^q$. Then B has a crystal structure, where the maps $\varepsilon_i,\ \varphi_i$ are given by

$$\varepsilon_i(b) = \max\{k \geq 0 \mid \tilde{e}_i b \neq 0\}, \qquad \varphi_i(b) = \max\{k \geq 0 \mid \tilde{f}_i b \neq 0\}.$$

In particular, we denote by $B(\lambda)$ the crystal of the irreducible highest weight module $V(\lambda)$ with highest weight $\lambda \in P^+$.

(2) Let $(L(\infty), B(\infty))$ be a crystal basis of $U_q^-(\mathfrak{g})$. Then $B(\infty)$ has a crystal structure, where the maps $\varepsilon_i,\ \varphi_i$ are given by

$$\varepsilon_i(b) = \max\{k \geq 0 \mid \tilde{e}_i b \neq 0\}, \qquad \varphi_i(b) = \varepsilon_i(b) + \langle h_i, \mathrm{wt}(b) \rangle.$$

(3) For $\lambda \in P$, let us consider $T_\lambda = \{t_\lambda\}$ with the maps:

$$\mathrm{wt}(t_\lambda) = \lambda, \quad \tilde{e}_i t_\lambda = \tilde{f}_i t_\lambda = 0 \text{ for } i \in I,$$
$$\varepsilon_i(t_\lambda) = \varphi_i(t_\lambda) = -\infty \text{ for } i \in I.$$

Then T_λ is a crystal.

(4) Let $C = \{c\}$. We define the maps

$$\mathrm{wt}(c) = 0, \quad \tilde{e}_i c = \tilde{f}_i c = 0, \quad \varepsilon_i(c) = \varphi_i(c) = 0 \ \ (i \in I).$$

Then C is a crystal.

Definition 1.4. Let B_1 and B_2 be crystals.

(1) A map $\psi : B_1 \to B_2$ is a *crystal morphism* if it satisfies the following properties:
  (a) for $b \in B_1$, we have

$$\mathrm{wt}(\psi(b)) = \mathrm{wt}(b),\ \varepsilon_i(\psi(b)) = \varepsilon_i(b),\ \varphi_i(\psi(b)) = \varphi_i(b) \text{ for all } i \in I,$$

  (b) for $b \in B_1$ and $i \in I$ with $\tilde{f}_i b \in B_1$, we have $\psi(\tilde{f}_i b) = \tilde{f}_i \psi(b)$.

(2) A crystal morphism $\psi : B_1 \to B_2$ is called *strict* if

$$\psi(\tilde{e}_i b) = \tilde{e}_i \psi(b),\ \psi(\tilde{f}_i b) = \tilde{f}_i \psi(b) \quad \text{for all } i \in I \text{ and } b \in B_1.$$

Here, we understand $\psi(0) = 0$.

(3) ψ is called an *embedding* if the underlying map $\psi : B_1 \to B_2$ is injective.

Let B_1 and B_2 be crystals. The *tensor product* $B_1 \otimes B_2$ is defined to be the set $B_1 \times B_2$ together with the following maps:

(a) $\mathrm{wt}(b_1 \otimes b_2) = \mathrm{wt}(b_1) + \mathrm{wt}(b_2)$,

(b) $\varepsilon_i(b_1 \otimes b_2) = \max\{\varepsilon_i(b_1),\ \varepsilon_i(b_2) - \langle h_i, \mathrm{wt}(b_1)\rangle\}$,

(c) $\varphi_i(b_1 \otimes b_2) = \max\{\varphi_i(b_2),\ \varphi_i(b_1) + \langle h_i, \mathrm{wt}(b_2)\rangle\}$,

(d) $\tilde{e}_i(b_1 \otimes b_2) = \begin{cases} \tilde{e}_i b_1 \otimes b_2 & \text{if } \varphi_i(b_1) \geq \varepsilon_i(b_2), \\ b_1 \otimes \tilde{e}_i b_2 & \text{if } \varphi_i(b_1) < \varepsilon_i(b_2), \end{cases}$

(e) $\tilde{f}_i(b_1 \otimes b_2) = \begin{cases} \tilde{f}_i b_1 \otimes b_2 & \text{if } \varphi_i(b_1) > \varepsilon_i(b_2), \\ b_1 \otimes \tilde{f}_i b_2 & \text{if } \varphi_i(b_1) \leq \varepsilon_i(b_2). \end{cases}$

It was shown in [11] that there is a unique strict crystal embedding

$$B(\lambda) \hookrightarrow B(\infty) \otimes T_\lambda \otimes C$$

sending u_λ to $\mathbf{1} \otimes t_\lambda \otimes c$. Here, u_λ is the highest weight element of $B(\lambda)$. We denote by ι_λ the composition of the strict embedding and the natural projection:

$$\iota_\lambda : B(\lambda) \hookrightarrow B(\infty) \otimes T_\lambda \otimes C \longrightarrow B(\infty). \tag{1.1}$$

Note that ι_λ is injective, but not a crystal morphism.

2. Path realization

Let $U_q'(\mathfrak{g})$ be the subalgebra of $U_q(\mathfrak{g})$ generated by $e_i, f_i, q^{\pm h_i}$ $(i \in I)$ and we set $\overline{P}^\vee = \bigoplus_{i=0}^n \mathbb{Z} h_i$, $\overline{\mathfrak{h}} = \mathbb{C} \otimes_{\mathbb{Z}} \overline{P}^\vee$, $\overline{P} = \bigoplus_{i=0}^n \mathbb{Z}\Lambda_i$ and $\overline{P}^+ = \sum_{i=0}^n \mathbb{Z}_{\geq 0}\Lambda_i$. Denote by $\mathrm{cl} : P \to \overline{P}$ the natural projection from P to $\overline{P}$. An abstract crystal B associated with $U_q'(\mathfrak{g})$ is called a *classical crystal.* For $b \in B$, we define

$$\varepsilon(b) = \sum_{i=0}^n \varepsilon_i(b)\Lambda_i, \qquad \varphi(b) = \sum_{i=0}^n \varphi_i(b)\Lambda_i.$$

Definition 2.1. A *perfect crystal of level* ℓ is a finite classical crystal B satisfying the following conditions:

(a) there exists a finite dimensional $U_q'(\mathfrak{g})$-module with a crystal basis whose crystal graph is isomorphic to B,
(b) $B \otimes B$ is connected,
(c) there exists a classical weight $\lambda_0 \in \overline{P}$ such that

$$\mathrm{wt}(B) \subset \lambda_0 + \sum_{i \neq 0} \mathbb{Z}_{\leq 0}\alpha_i, \qquad \#(B_{\lambda_0}) = 1,$$

(d) for any $b \in B$, we have $\langle c,\ \varepsilon(b)\rangle \geq \ell$,
(e) for each $\lambda \in \overline{P}_\ell^+ := \{\mu \in \overline{P}^+ \mid \langle c,\ \mu\rangle = \ell\}$, there exist unique vectors b^λ and b_λ in B such that $\varepsilon(b^\lambda) = \lambda,\ \varphi(b_\lambda) = \lambda$.

Given a dominant integral weight λ with $\lambda(c) = \ell$ and a perfect crystal B of level ℓ, it was shown in [7] that there exists a unique crystal isomorphism, called the *fundamental isomorphism theorem for perfect crystals*,

$$\psi : B(\lambda) \xrightarrow{\sim} B(\varepsilon(b_\lambda)) \otimes B \tag{2.1}$$

sending u_λ to $u_{\varepsilon(b_\lambda)} \otimes b_\lambda$. By applying this crystal isomorphism repeatedly, we get a sequence of crystal isomorphisms

$$B(\lambda) \xrightarrow{\sim} B(\lambda_1) \otimes B \xrightarrow{\sim} B(\lambda_2) \otimes B \otimes B \xrightarrow{\sim} \cdots\cdots,$$

where $\lambda_0 = \lambda$, $b_0 = b_\lambda$, $\lambda_{k+1} = \varepsilon(b_k)$, $b_{k+1} = b_{\lambda_{k+1}}$ $(k \geq 0)$. The sequence $\mathbf{p}_\lambda := (b_k)_{k=0}^\infty$ is called the *ground-state path of weight* λ and a sequence $\mathbf{p} = (p_k)_{k=0}^\infty$ of elements $p_k \in B$ is called a λ*-path in* B if $p_k = b_k$ for all $k \gg 0$. We denote by $\mathcal{P}^B(\lambda)$ the set of λ-paths in B, which gives rise to the *path realization* of $B(\lambda)$.

Theorem 2.2 ([7]). *There exists a unique crystal isomorphism* $B(\lambda) \xrightarrow{\sim} \mathcal{P}^B(\lambda)$ *which maps* u_λ *to* $\mathbf{p}_\lambda$.

We list some examples of perfect crystals of level 1 and the corresponding ground-state paths (see [1, 8], etc).

Example 2.3.

(1) The crystal B^1 and its ground-state $\mathbf{p}^1_{\Lambda_i}$ of weight Λ_i $(i \in I)$ are given as

$$\begin{tikzcd} \mathbf{b}_1 \arrow[r, "1"] & \mathbf{b}_2 \arrow[r, "2"] & \cdots \arrow[r, "n-1"] & \mathbf{b}_n \arrow[r, "n"] & \mathbf{b}_{n+1} \arrow[llll, bend left, "0"] \end{tikzcd},$$

$$\mathbf{p}^1_{\Lambda_i} = (\cdots, \mathbf{b}_1, \mathbf{b}_2, \cdots, \mathbf{b}_n, \mathbf{b}_{n+1}, \mathbf{b}_1, \mathbf{b}_2, \cdots, \mathbf{b}_{i-1}, \mathbf{b}_i).$$

We denote by $\mathcal{P}^1(\Lambda_i)$ the set of all Λ_i-paths in B^1.

(2) The crystal B^n and its ground-state $\mathbf{p}^n_{\Lambda_i}$ of weight Λ_i $(i \in I)$ are given as

$$\begin{tikzcd} \overline{\mathbf{b}}_{n+1} \arrow[r, "n"] & \overline{\mathbf{b}}_n \arrow[r, "n-1"] & \cdots \arrow[r, "2"] & \overline{\mathbf{b}}_2 \arrow[r, "1"] & \overline{\mathbf{b}}_1 \arrow[llll, bend left, "0"] \end{tikzcd},$$

$$\mathbf{p}^n_{\Lambda_i} = (\cdots, \overline{\mathbf{b}}_{n+1}, \overline{\mathbf{b}}_n, \cdots, \overline{\mathbf{b}}_2, \overline{\mathbf{b}}_1, \overline{\mathbf{b}}_{n+1}, \overline{\mathbf{b}}_n, \cdots, \overline{\mathbf{b}}_{i+2}, \overline{\mathbf{b}}_{i+1}).$$

We denote by $\mathcal{P}^n(\Lambda_i)$ the set of all Λ_i-paths in B^n.

(3) The *adjoint crystal* B^{ad} is given as follows.

Let

$$B^{\mathrm{ad}} = \{\emptyset\} \cup \{\mathsf{b}_{\pm\alpha_{ij}} \mid 1 \le i \le j \le n\} \cup \{h_i \mid i = 1, \ldots, n\},$$

where $\alpha_{ij} := \alpha_i + \alpha_{i+1} + \cdots + \alpha_j$ for $1 \le i \le j \le n$. We define the i-arrows $(i \in I)$ by

$$\begin{aligned} (i \neq 0) \quad & \mathsf{b}_\alpha \xrightarrow{i} \mathsf{b}_\beta \ \Leftrightarrow \ \alpha - \alpha_i = \beta, \\ & \mathsf{b}_{\alpha_i} \xrightarrow{i} h_i \xrightarrow{i} \mathsf{b}_{-\alpha_i}, \\ (i = 0) \quad & \mathsf{b}_\alpha \xrightarrow{0} \mathsf{b}_\beta \ \Leftrightarrow \ \alpha + \theta = \beta \ (\alpha, \beta \neq \pm\theta), \\ & \mathsf{b}_{-\theta} \xrightarrow{0} \emptyset \xrightarrow{0} \mathsf{b}_\theta, \end{aligned}$$

where $\theta = \alpha_1 + \cdots + \alpha_n$. The crystal B^{ad} is a perfect crystal of level 1 with the ground-state path of weight Λ_0

$$\mathbf{p}^{\mathrm{ad}}_{\Lambda_0} = (\cdots, \emptyset, \emptyset, \cdots, \emptyset, \emptyset).$$

There is a crystal isomorphism $\mathfrak{p}^{\mathrm{ad}} : B^n \otimes B^1 \to B^{\mathrm{ad}}$ given by

$$\mathfrak{p}^{\mathrm{ad}}(\overline{\mathbf{b}}_j \otimes \mathbf{b}_i) = \begin{cases} \mathsf{b}_{\mathrm{wt}(\overline{\mathbf{b}}_j \otimes \mathbf{b}_i)} & \text{if } \mathrm{wt}(\overline{\mathbf{b}}_j \otimes \mathbf{b}_i) \neq 0, \\ h_i & \text{if } \mathrm{wt}(\overline{\mathbf{b}}_j \otimes \mathbf{b}_i) = 0, i \neq n+1, \\ \emptyset & \text{otherwise}. \end{cases} \tag{2.2}$$

We denote by $\mathcal{P}^{\mathrm{ad}}(\Lambda_0)$ the set of all Λ_0-paths in B^{ad}.

3. Combinatorics of Young Walls

In [4, 6], Kang gave a combinatorial realization of crystal graphs for basic representations of quantum affine algebras of type $A_n^{(1)}$ $(n \geq 1)$, $A_{2n-1}^{(2)}$ $(n \geq 3)$, $D_n^{(1)}$ $(n \geq 4)$, $A_{2n}^{(2)}$, $D_{n+1}^{(2)}$ $(n \geq 2)$, and $B_n^{(1)}$ $(n \geq 3)$ by using new combinatorial objects called *Young walls*, which are a generalization of colored Young diagrams used in [2, 5, 15]. In this work, we focus on the quantum affine algebra $U_q(A_n^{(1)})$.

The *Young wall* Y^1 (resp. Y^n) is a wall consisting of colored blocks stacked by the following rules:

(a) the colored blocks should be stacked in the pattern P^1 (resp. P^n) of weight Λ_k given below,

(b) except for the right-most column, there should be no free space to the right of any block.

The patterns are given as follows.



the pattern P^1 of weight Λ_k



the pattern P^n of weight Λ_k

Note that the heights of the columns of a Young wall Y are weakly decreasing from right to left, so we denote it by $Y = (y_i)_{i \geq 0}$, where y_i is the i-th column of Y.

Definition 3.1. Let Y be a Young wall corresponding to the pattern P^1 (resp. P^n).

(1) An i-block in Y is called a *removable i-block* if Y remains a Young wall after removing the block.

(2) A place in Y is called an *admissible i-slot* if one may add an i-block to obtain another Young wall.

(3) A column in Y is said to be *i-removable* (resp. *i-admissible*) if the column has a removable i-block (resp. an admissible i-slot).

Now we define the action of Kashiwara operators $\tilde{e}_i, \tilde{f}_i$ $(i \in I)$ on Young walls. Let $Y = (y_k)_{k \geq 0}$ be a Young wall corresponding to the pattern P^1 (resp. P^n).

(a) To each column y_k of Y, we assign

$$\begin{cases} - & \text{if } y_k \text{ is } i\text{-removable}, \\ + & \text{if } y_k \text{ is } i\text{-admissible}, \end{cases}$$

(b) From this sequence of $+$'s and $-$'s, cancel out all $(+,-)$ pairs to obtain a finite sequence of $-$'s followed by $+$'s. This sequence $(-,\cdots,-,+,\cdots,+)$ is called the *i-signature* of Y.

(c) We define $\tilde{e}_iY$ to be the Young wall obtained from Y by removing the i-block corresponding to the rightmost $-$ in the i-signature of Y. If there is no $-$ in the i-signature, we set $\tilde{e}_iY = 0$.

(d) We define $\tilde{f}_iY$ to be the Young wall obtained from Y by adding an i-block to the column corresponding to the leftmost $+$ in the i-signature of Y. If there is no $+$ in the i-signature, we set $\tilde{f}_iY = 0$.

We also define

$$\begin{aligned}
\mathrm{wt}(y_j) &= \sum_{i\in I} k_{ij}\alpha_i \ (j \in \mathbb{Z}_{\geq 0}),\\
\mathrm{wt}(Y) &= \Lambda_k - \sum_{j\geq 0}\mathrm{wt}(y_j),\\
\varepsilon_i(Y) &= \text{ the number of }-\text{'s in the }i\text{-signature of }Y,\\
\varphi_i(Y) &= \text{ the number of }+\text{'s in the }i\text{-signature of }Y,
\end{aligned}$$

where k_{ij} is the number of i-blocks in the j-th column y_j of Y. Note that the height of y_j is $\mathrm{ht}(\mathrm{wt}(y_j))$. Let $\mathcal{Y}^1(\Lambda_k)$ (resp. $\mathcal{Y}^n(\Lambda_k)$) be the set of all Young walls Y^1 (resp. Y^n) whose shapes are $(n+1)$-reduced Young diagrams; i.e., $Y^1 = (y_j)_{j\geq 0} \in \mathcal{Y}^1(\Lambda_k)$ (resp. $Y^n = (\overline{y}_j)_{j\geq 0} \in \mathcal{Y}^n(\Lambda_k)$) if and only if

$$\mathrm{ht}(\mathrm{wt}(y_j)) - \mathrm{ht}(\mathrm{wt}(y_{j+1})) < n+1 \quad (\text{resp. } \mathrm{ht}(\mathrm{wt}(\overline{y}_j)) - \mathrm{ht}(\mathrm{wt}(\overline{y}_{j+1})) < n+1)$$

for $j \geq 0$. Then $\mathcal{Y}^1(\Lambda_k)$ (resp. $\mathcal{Y}^n(\Lambda_k)$) has a $U_q(\mathfrak{g})$-crystal structure, and we have the following theorem.

Theorem 3.2. [4, 6, 15] *There is a unique crystal isomorphism* $B(\Lambda_k) \xrightarrow{\sim} \mathcal{Y}^1(\Lambda_k)$ (*resp.* $\mathcal{Y}^n(\Lambda_k)$) *which maps the highest weight element* u_{Λ_k} *to the empty Young wall* $\emptyset$.

Let $\mathbf{p}^1 = (\cdots,\mathbf{b}_{i_2},\mathbf{b}_{i_1},\mathbf{b}_{i_0})$ be a Λ_k-path in B^1. Consider a Young wall $\mathbf{Y}^1_k(\mathbf{p}^1) = (y_j(\mathbf{p}^1))_{j\geq 0}$ such that the j-th column $y_j(\mathbf{p}^1)$ is $\emptyset$ if $j > \mathrm{ht}(\Lambda_k - \mathrm{wt}(\mathbf{p}^1))$, otherwise $y_j(\mathbf{p}^1)$ is the smallest j-th column in P^1 satisfying the following conditions:

(a) the top color of $y_j(\mathbf{p}^1)$ is $i_j - 1$,

(b) $y_{j+1}(\mathbf{p}^1) \leq y_j(\mathbf{p}^1)$.

One can prove that the Young wall $\mathbf{Y}^1_k(\mathbf{p}^1)$ is contained in $\mathcal{Y}^1(\Lambda_k)$, and the map

$$\mathbf{Y}^1_k : \mathcal{P}^1(\Lambda_k) \longrightarrow \mathcal{Y}^1(\Lambda_k) \tag{3.1}$$

is a crystal isomorphism. If we set $Y^1 = (y_j)_{j\geq 0} \in \mathcal{Y}^1(\Lambda_k)$, then the inverse image $\mathbf{p}^1$ of Y^1 under the crystal isomorphism $\mathbf{Y}^1_k$ is

$$\mathbf{p}^1 = (\ldots,\mathbf{b}_{a_j},\ldots,\mathbf{b}_{a_1},\mathbf{b}_{a_0}), \tag{3.2}$$

where $a_j \equiv \mathrm{ht}(\mathrm{wt}(y_j)) - j + k \pmod{n+1}$ for all $j \geq 0$.

In a similar manner, given a Λ_k-path $\mathbf{p}^n = (\cdots, \overline{\mathbf{b}}_{i_2}, \overline{\mathbf{b}}_{i_1}, \overline{\mathbf{b}}_{i_0})$ in B^n, we have a Young wall $\mathbf{Y}_k^n(\mathbf{p}^n) = (\overline{y}_j(\mathbf{p}^n))_{j\geq 0}$ such that the j-th column $\overline{y}_j(\mathbf{p}^n)$ is $\emptyset$ if $j > \mathrm{ht}(\Lambda_k - \mathrm{wt}(\mathbf{p}^n))$, otherwise $\overline{y}_j(\mathbf{p}^n)$ is the smallest j-th column in P^n satisfying the following conditions:

(a) the top color of $\overline{y}_j(\mathbf{p}^n)$ is i_j,
(b) $\overline{y}_{j+1}(\mathbf{p}^n) \leq \overline{y}_j(\mathbf{p}^n)$.

One can prove that the Young wall $\mathbf{Y}_k^n(\mathbf{p}^n)$ is contained in $\mathcal{Y}^n(\Lambda_k)$, and the map

$$\mathbf{Y}_k^n : \mathcal{P}^n(\Lambda_k) \longrightarrow \mathcal{Y}^n(\Lambda_k) \tag{3.3}$$

is a crystal isomorphism. If we set $Y^n = (\overline{y}_j)_{j\geq 0} \in \mathcal{Y}^n(\Lambda_k)$, then the inverse image $\mathbf{p}^n$ of Y^n under the crystal isomorphism $\mathbf{Y}_k^n$ is

$$\mathbf{p}^n = (\ldots, \overline{\mathbf{b}}_{b_j}, \ldots, \overline{\mathbf{b}}_{b_1}, \overline{\mathbf{b}}_{b_0}), \tag{3.4}$$

where $b_j \equiv 1 - \mathrm{ht}(\mathrm{wt}(\overline{y}_j)) + j + k \pmod{n+1}$ for all $j \geq 0$.

Example 3.3. Let $\mathfrak{g}$ be the Kac-Moody algebra of type $A_3^{(1)}$. Fix an element

$$b = \tilde{f}_0\tilde{f}_2\tilde{f}_1(\tilde{f}_1\tilde{f}_2\tilde{f}_3\tilde{f}_0)^3 u_{\Lambda_0} \in B(\Lambda_0),$$

where u_{Λ_0} is the highest weight element of $B(\Lambda_0)$. Then the Λ_0-paths $\mathbf{p}^1 \in \mathcal{P}^1(\Lambda_0)$, $\mathbf{p}^n \in \mathcal{P}^n(\Lambda_0)$ and $\mathbf{p}^{\mathrm{ad}} \in \mathcal{P}^{\mathrm{ad}}(\Lambda_0)$ corresponding to b are given as

$$\begin{aligned}
\mathbf{p}^1 &= (\ldots, \mathbf{b}_3, \mathbf{b}_1, \mathbf{b}_2, \mathbf{b}_3, \mathbf{b}_0, \mathbf{b}_1, \mathbf{b}_2, \mathbf{b}_3, \mathbf{b}_0, \mathbf{b}_1, \mathbf{b}_2, \mathbf{b}_3, \mathbf{b}_0, \mathbf{b}_3),\\
\mathbf{p}^n &= (\ldots, \overline{\mathbf{b}}_3, \overline{\mathbf{b}}_2, \overline{\mathbf{b}}_0, \overline{\mathbf{b}}_1, \overline{\mathbf{b}}_0, \overline{\mathbf{b}}_2, \overline{\mathbf{b}}_1),\\
\mathbf{p}^{\mathrm{ad}} &= (\ldots, \emptyset, \emptyset, \mathsf{b}_{\alpha_1+\alpha_2+\alpha_3}, h_1, h_1, \mathsf{b}_{-\alpha_1-\alpha_2}),
\end{aligned}$$

which yield the Young walls $Y^1 := \mathbf{Y}_0^1(\mathbf{p}^1)$ and $Y^n := \mathbf{Y}_0^n(\mathbf{p}^n)$ as follows:

$$Y^1 = \begin{array}{|c|c|c|c|c|c|c|c|c|c|c|c|c|}
\multicolumn{12}{c|}{} & 2 \\ \cline{13-13}
\multicolumn{12}{c|}{} & 1 \\ \hline
0 & 1 & 2 & 3 & 0 & 1 & 2 & 3 & 0 & 1 & 2 & 3 & 0 \\ \hline
\end{array}\,, \qquad
Y^n = \begin{array}{|c|c|c|c|c|}
\cline{4-5}
\multicolumn{3}{c|}{} & 2 & 1 \\ \cline{2-5}
\multicolumn{1}{c|}{} & 1 & 0 & 3 & 2 \\ \cline{2-5}
\multicolumn{1}{c|}{} & 2 & 1 & 0 & 3 \\ \hline
0 & 3 & 2 & 1 & 0 \\ \hline
\end{array}\,.$$

4. Geometric Constructions of Crystal Graphs

In this section, we review geometric constructions of crystal bases via quiver varieties. See [3, 12, 14, 17, 18, 19] for more details.

Let $I = \mathbb{Z}/(n+1)\mathbb{Z}$ and H the set of the arrows such that $i \to j$ with $i, j \in I$, $i - j = \pm 1$. For $h \in H$, we denote by $\mathrm{in}(h)$ (resp. $\mathrm{out}(h)$) the incoming (resp. outgoing) vertex of h. Define an involution $- : H \to H$ to be the map interchanging $i \to j$ and $j \to i$. Let

$$\Omega = \{h \in H \mid \mathrm{in}(h) - \mathrm{out}(h) = 1\}$$

so that $H = \Omega \sqcup \overline{\Omega}$; i.e.,

$(I,\Omega) =$  ,

$(I,\overline{\Omega}) =$  .

Note that our graph is an affine Dynkin graph of type $A_n^{(1)}$. We take the map $\epsilon : H \to \{-1, 1\}$ given by

$$\epsilon(h) = \begin{cases} 1 & \text{if } h \in \Omega, \\ -1 & \text{if } h \in \overline{\Omega}. \end{cases}$$

For $\alpha = \sum_{i=0}^{n} k_i\alpha_i \in Q^+$, we define the I-graded vector space

$$V(\alpha) = \bigoplus_{i=0}^{n} V_i(\alpha),$$

where $V_i(\alpha)$ is the $\mathbb{C}$-vector space with an ordered basis $v^i(\alpha) = \{v_0^i, v_1^i \ldots, v_{k_i-1}^i\}$ for all i. Fix an ordered basis

$$v(\alpha) = \{v_0^0, \ldots, v_{k_0-1}^0, v_0^1, \ldots, v_{k_1-1}^1, \ldots, v_0^n, \ldots v_{k_n-1}^n\},$$

for $V(\alpha)$ and set

$$\underline{\dim} V(\alpha) = \sum_{i=0}^{n} k_i\alpha_i = \alpha.$$

In a similar manner, for $\lambda = \sum_{i=0}^{n} w_i\Lambda_i \in P^+$, we define the I-graded vector space

$$W(\lambda) = \bigoplus_{i=0}^{n} W_i(\lambda),$$

where $W_i(\lambda)$ is a $\mathbb{C}$-vector space of dimension w_i.

Given $\alpha \in Q^+$, we set $V = V(\alpha)$ (resp. $V_i = V_i(\alpha)$ $(i = I)$) and let

$$E(\alpha) = E_\Omega(\alpha) \oplus E_{\overline{\Omega}}(\alpha),$$

where

$$E_\Omega(\alpha) = \bigoplus_{h\in\Omega} \mathrm{Hom}(V_{\mathrm{out}(h)}, V_{\mathrm{in}(h)}) = \bigoplus_{i\in I} \mathrm{Hom}(V_{i-1},\ V_i),$$
$$E_{\overline{\Omega}}(\alpha) = \bigoplus_{h\in\overline{\Omega}} \mathrm{Hom}(V_{\mathrm{out}(h)}, V_{\mathrm{in}(h)}) = \bigoplus_{i\in I} \mathrm{Hom}(V_i,\ V_{i-1}).$$

Let us denote by π_Ω (resp. $\pi_{\overline{\Omega}}$) the natural projection from $E(\alpha)$ to $E_\Omega(\alpha)$ (resp. $E_{\overline{\Omega}}(\alpha)$). For $\chi \in E(\alpha)$, if there is no danger of confusion, we write $x = (x_i \in \mathrm{Hom}(V_{i-1},\ V_i))_{i\in I}$ (resp. $\overline{x} = (\overline{x}_i \in \mathrm{Hom}(V_i,\ V_{i-1}))_{i\in I}$) for $\pi_\Omega(\chi)$ (resp. $\pi_{\overline{\Omega}}(\chi)$). The matrix representation of $x \in E_\Omega(\alpha)$ in the ordered basis $v(\alpha)$ is given as

$$x = \begin{pmatrix} 0 & & \cdots & 0 & x_0 \\ x_1 & 0 & & & 0 \\ 0 & x_2 & 0 & & \vdots \\ \vdots & & \ddots & \ddots & 0 \\ 0 & \cdots & 0 & x_n & 0 \end{pmatrix},$$

where x_i is the matrix representation of $x|_{V_{i-1}} : V_{i-1} \to V_i$ in the ordered bases $v^{i-1}(\alpha)$ and $v^i(\alpha)$. We also may consider the matrix representation of $\overline{x} \in E_{\overline{\Omega}}(\alpha)$ in the same manner:

$$\overline{x} = \begin{pmatrix} 0 & \overline{x}_1 & 0 & \cdots & 0 \\ & 0 & \overline{x}_2 & & \vdots \\ \vdots & & 0 & \ddots & 0 \\ 0 & & & \ddots & \overline{x}_n \\ \overline{x}_0 & 0 & \cdots & 0 & 0 \end{pmatrix},$$

where $\overline{x}_i$ is the matrix representation of $\overline{x}|_{V_i} : V_i \to V_{i-1}$ in the ordered bases $v^i(\alpha)$ and $v^{i-1}(\alpha)$.

For $s \in I,\ 0 \le i < k_{s-1},\ 0 \le j < k_s$, define the linear map $\mathcal{E}^s_{ij} : V_{s-1} \to V_s$ by

$$\mathcal{E}^s_{ij}(v^{s-1}_k) = \begin{cases} v^s_j & \text{if } k = i, \\ 0 & \text{otherwise.} \end{cases}$$

Similarly, for $s \in I,\ 0 \le i < k_s,\ 0 \le j < k_{s-1}$, define the linear map $\overline{\mathcal{E}}^s_{ij} : V_s \to V_{s-1}$ by

$$\overline{\mathcal{E}}^s_{ij}(v^s_k) = \begin{cases} v^{s-1}_j & \text{if } k = i, \\ 0 & \text{otherwise.} \end{cases}$$

The algebraic group $G(\alpha) := \prod_{i \in I} \mathrm{Aut}(V_i) \subset \mathrm{Aut}(V)$ acts on $E(\alpha)$ by $(g, \chi) = g\chi g^{-1}$ for $g \in G(\alpha),\ \chi \in E(\alpha)$. Let $\langle \cdot, \cdot \rangle$ be the nondegenerate, $G(\alpha)$-invariant, sympletic form on $E(\alpha)$ defined by

$$\langle \chi, \chi' \rangle = \sum_{h \in H} \epsilon(h) \mathrm{tr}(\chi_h \chi'_{\overline{h}})$$

for $\chi, \chi' \in E(\alpha)$. Note that $E(\alpha)$ may be viewed as the cotangent bundle of $E_\Omega(\alpha)$ (resp. $E_{\overline{\Omega}}(\alpha)$) under this form. The *moment map* $\mu = (\mu_i : E(\alpha) \to \mathrm{End}(V_i))_{i \in I}$ is given by

$$\mu_i(\chi) = \sum_{h \in H,\ \mathrm{in}(h) = i} \epsilon(h) \chi_h \chi_{\overline{h}} \ = \ x_i \overline{x}_i - \ \overline{x}_{i+1} x_{i+1}$$

for $\chi = x + \overline{x} \in E(\alpha)$. Note that

$$\mu_i(\chi) = 0 \text{ for all } i \in I \quad \text{if and only if} \quad [x, \overline{x}] = x\overline{x} - \overline{x}x = 0. \tag{4.1}$$

Here, $x\overline{x} = (x_i \overline{x}_i : V_i \to V_i)_{i \in I}$ and $\overline{x}x = (\overline{x}_{i+1} x_{i+1} : V_i \to V_i)_{i \in I}$.

An element $\chi \in E(\alpha)$ is *nilpotent* if there exists an $N \geq 2$ such that for any sequence $h_1, \dots, h_N \in H$ satisfying $\mathrm{in}(h_i) = \mathrm{out}(h_{i+1})$ $(i = 1, \dots, N-1)$, the composition map $\chi_{h_N} \cdots \chi_{h_1}$ is zero. We define *Lusztig's quiver variety* to be

$$\Lambda(\alpha) = \{\chi \in E(\alpha) |\ \chi : \text{nilpotent},\ \mu_i(\chi) = 0 \text{ for all } i \in I\}.$$

We denote by $\mathrm{Irr}\Lambda(\alpha)$ the set of all irreducible components of $\Lambda(\alpha)$

For a pair $(k' \leq k)$ of integers, let $V(k', k) = \bigoplus_{i \in I} V_i(k', k)$ be the I-graded vector space with basis $\{\mathsf{e}_j |\ k' \leq j \leq k\}$ such that $V_i(k', k) = \mathrm{span}_{\mathbb{C}}\{\mathsf{e}_j |\ j \equiv i \pmod{n+1}\}$ for $i \in I$. Consider the $\mathbb{C}$-linear map $x(k', k) : V(k', k) \to V(k', k)$ sending e_i to e_{i+1}, where $\mathsf{e}_{k+1} = 0$. Then it is clear that the representation $(V(k', k), x(k', k))$ of the quiver (I, Ω) is indecomposable and nilpotent. Note that the isomorphism class of $(V(k', k), x(k', k))$ does not change when k' and k are simultaneously translated by a multiple $n+1$. Moreover, any indecomposable nilpotent finite-dimensional representation of the quiver (I, Ω) is isomorphic to $(V(k', k), x(k', k))$ for some pair $(k' \leq k)$. Let $\mathbf{Z}$ be the set of all pairs $(k' \leq k)$ of integers defined up to simultaneous translation by a multiple of $n+1$ and let $\tilde{\mathbf{Z}}$ be the set of all functions from $\mathbf{Z}$ to $\mathbb{Z}_{\geq 0}$ with finite support. Note that $\tilde{\mathbf{Z}}$ naturally corresponds to isomorphism classes of nilpotent finite-dimensional representations of the quiver (I, Ω). The set of $G(\alpha)$-orbits on the set of nilpotent elements in $E_\Omega(\alpha)$ is naturally indexed by the subset $\tilde{\mathbf{Z}}(\alpha)$ of $\tilde{\mathbf{Z}}$ such that, for $f \in \tilde{\mathbf{Z}}(\alpha)$,

$$\sum_{k' \leq k} f(k', k) \cdot \#\{j |\ k' \leq j \leq k,\ j \equiv i \pmod{n+1}\} = \dim V_i \ (i \in I).$$

Here the sum is taken over all $k' \leq k$ up to simultaneous translation by a multiple of $n+1$. An element $f \in \tilde{\mathbf{Z}}(\alpha)$ is *aperiodic* if, for any $k' \leq k$, not all integers $f(k', k), f(k'+1, k+1), \dots, f(k'+n, k+n)$ are greater than zero. For any $f \in \tilde{\mathbf{Z}}(\alpha)$, let $\mathcal{C}_f$ be the conormal bundle of the $G(\alpha)$-orbit corresponding to f, and let $\overline{\mathcal{C}}_f$ be the closure of $\mathcal{C}_f$. Then we have

Theorem 4.1 ([14])**.** *The map $f \mapsto \overline{\mathcal{C}}_f$ is a 1-1 correspondence between the set of aperiodic elements in $\tilde{\mathbf{Z}}(\alpha)$ and* $\mathrm{Irr}\Lambda(\alpha)$.

In a similar manner, for a pair $(k \geq k')$ of integers, let $\overline{x}(k, k') : V(k', k) \to V(k', k)$ be the $\mathbb{C}$-linear map sending e_i to e_{i-1}, where $\mathsf{e}_{k'-1} = 0$. Then the representation $(V(k', k), \overline{x}(k, k'))$ of the quiver $(I, \overline{\Omega})$ is indecomposable and nilpotent, and the isomorphism class of $(V(k', k), \overline{x}(k, k'))$ does not change when k and k' are simultaneously translated by a multiple $n+1$. Any indecomposable nilpotent finite-dimensional representation of the quiver $(I, \overline{\Omega})$ is isomorphic to $(V(k', k), \overline{x}(k, k'))$ for some pair $(k \geq k')$. Let $\overline{\mathbf{Z}}$ be the set of all pairs $(k \geq k')$ of integers defined up to simultaneous translation by a multiple of $n+1$ and let $\tilde{\overline{\mathbf{Z}}}$ be the set of all functions from $\overline{\mathbf{Z}}$ to $\mathbb{Z}_{\geq 0}$ with finite support. Then the set of $G(\alpha)$-orbits on the set of nilpotent elements in $E_{\overline{\Omega}}(\alpha)$ is naturally indexed by the subset $\tilde{\overline{\mathbf{Z}}}(\alpha)$ of $\tilde{\overline{\mathbf{Z}}}$ such that, for $f \in \tilde{\overline{\mathbf{Z}}}(\alpha)$,

$$\sum_{k \geq k'} f(k, k') \cdot \#\{j |\ k \geq j \geq k',\ j \equiv i \pmod{n+1}\} = \dim V_i \ (i \in I).$$

Here the sum is taken over all $k \geq k'$ up to simultaneous translation by a multiple of $n+1$. An element $f \in \tilde{\overline{\mathbf{Z}}}(\alpha)$ is *aperiodic* if, for any $k \geq k'$, not all integers $f(k, k'), f(k+1, k'+1), \dots, f(k+n, k'+n)$ are greater than zero. Then one can show that there is a 1-1 correspondence between the set of aperiodic elements in $\tilde{\tilde{\mathbf{Z}}}(\alpha)$ and $\mathrm{Irr}\Lambda(\alpha)$.

Moreover, Kashiwara and Saito [12] gave a crystal structure on

$$\mathbb{B}(\infty) := \bigsqcup_{\alpha \in Q^+} \mathrm{Irr}\Lambda(\alpha),$$

and proved the following theorem.

Theorem 4.2 ([12])**.** *There is a unique crystal isomorphism* $\mathbb{B}(\infty) \cong B(\infty)$.

Now we introduce a description of Nakajima's quiver varieties presented in [16]. Given $\alpha \in Q^+$ and $\lambda \in P^+$, we set $W = W(\lambda)$ (resp. $W_i = W_i(\lambda)$) and let

$$E(\lambda, \alpha) = \Lambda(\alpha) \times \sum_{i \in I} \mathrm{Hom}(V_i, W_i).$$

The group $G(\alpha)$ acts on $E(\lambda, \alpha)$ by $(g, (\chi, \mathfrak{t})) = (g\chi g^{-1}, \mathfrak{t}g^{-1})$. For $\chi \in \Lambda(\alpha)$, an I-graded subspace S of $V(\alpha)$ is *χ-stable* if $\chi_h(S_{\mathrm{out}(h)}) \subset S_{\mathrm{in}(h)}$ for all $h \in H$. An element $(\chi, \mathfrak{t}) \in E(\lambda, \alpha)$ is called a *stable point* of $E(\lambda, \alpha)$ if it satisfies the following conditions: if S is a χ-stable subspace of V with $\mathfrak{t}_i(S_i) = 0$ $(i \in I)$, then $S = 0$. Let $E(\lambda, \alpha)^{st}$ be the set of all stable points of $E(\lambda, \alpha)$, and define

$$\Lambda(\lambda, \alpha) = E(\lambda, \alpha)^{st}/G(\alpha).$$

Let $\mathrm{Irr}\Lambda(\lambda, \alpha)$ (resp. $\mathrm{Irr}E(\lambda, \alpha)$) be the set of all irreducible components of $\Lambda(\lambda, \alpha)$ (resp. $E(\lambda, \alpha)$). Since $\mathrm{Irr}\Lambda(\lambda, \alpha)$ can be identified with

$$\{Z \in \mathrm{Irr}E(\lambda, \alpha) |\ Z \cap E(\lambda, \alpha)^{st} \neq \emptyset\},$$

each irreducible component X in $\mathrm{Irr}\Lambda(\lambda, \alpha)$ can be written as

$$X = \left(\left(X_0 \times \sum_{i \in I} \mathrm{Hom}(V_i, W_i) \right) \cap E(\lambda, \alpha)^{st} \right) /G(\alpha)$$

for some irreducible component X_0 in $\mathrm{Irr}\Lambda(\alpha)$.

In [18], Saito gave a crystal structure on

$$\mathbb{B}(\lambda) := \bigsqcup_{\alpha \in Q^+} \mathrm{Irr}\Lambda(\lambda, \alpha),$$

and proved the following theorem.

Theorem 4.3 ([18])**.** *There is a unique crystal isomorphism* $\mathbb{B}(\lambda) \cong B(\lambda)$.

In [3], Frenkel and Savage gave an enumeration of $\mathrm{Irr}\Lambda(\lambda, \alpha)$ in terms of Young and Maya diagrams for type $A_n^{(1)}$. Combining Theorem 4.2 and Theorem 4.3 with (1.1), we obtain an injective map

$$\iota_\lambda : \mathbb{B}(\lambda) \hookrightarrow \mathbb{B}(\infty). \tag{4.2}$$

For each irreducible component $X_0 \in \iota_{\Lambda_k}(\mathbb{B}(\Lambda_k))$, Frenkel and Savage constructed a special point in $X_0 \times \sum_{i \in I} \mathrm{Hom}(V_i, W_i)$ which is not killed by the stability condition, and showed that there is a 1-1 correspondence between the set of such special points and the set of $(n+1)$-reduced colored Young diagrams. Savage later established a crystal isomorphism between $\mathbb{B}(\lambda)$ and Young walls for quantum affine algebras of type $A_n^{(1)}$ and $D_n^{(1)}$ in [19].

We briefly recall the result of [3] for type $A_n^{(1)}$ in terms of Young walls. Note that the orientation appeared in [3] is $\overline{\Omega}$. Take a Young wall $Y^n \in \mathcal{Y}^n(\Lambda_k)$ such that $\mathrm{wt}(Y^n) = \alpha$. Let l_i be the length of the i-th row of the Young wall Y^n $(i \geq 1)$ and let N be the height of Y^n. Set

$$A_{Y^n} := \{(l_i - i + k, 1 - i + k) \mid 1 \leq i \leq N\} \subset \overline{\mathbf{Z}}$$

and consider the function $f \in \tilde{\overline{\mathbf{Z}}}(\alpha)$ given by

$$f(s, s') = \begin{cases} 1 & \text{if } (s, s') \in A_{Y^n}, \\ 0 & \text{otherwise.} \end{cases}$$

Note that f is aperiodic. Let $\overline{\mathcal{C}}_f$ be the closure of the conormal bundle of the $G(\alpha)$-orbit $\mathcal{O}_f$ in $E_{\overline{\Omega}}(\alpha)$ corresponding to f, and define the irreducible component

$$X_{Y^n} := \left(\left(\overline{\mathcal{C}}_f \times \sum_{i \in I} \mathrm{Hom}(V_i, W_i) \right) \cap E(\Lambda_k, \alpha)^{st} \right) / G(\alpha) \in \mathrm{Irr}\Lambda(\Lambda_k, \alpha).$$

By [3, Theorem 5.5], the map $Y^n \mapsto X_{Y^n}$ is a 1-1 correspondence between

$$\{Y^n \in \mathcal{Y}^n(\Lambda_k) |\ \mathrm{wt}(Y^n) = \alpha\} \ \text{ and } \ \mathrm{Irr}\Lambda(\Lambda_k, \alpha).$$

Moreover, it is proved in [19, Theorem 8.4.] that the map $Y^n \mapsto X_{Y^n}$ from $\mathcal{Y}^n(\Lambda_k)$ to $\mathbb{B}(\Lambda_k)$ is a crystal isomorphism. We would like to point out that $\overline{\mathcal{C}}_f = \iota_{\Lambda_k}(X_{Y^n})$.

Now we construct an element in the $G(\alpha)$-orbit $\mathcal{O}_f$ in $E_{\overline{\Omega}}(\alpha)$ from the Young wall Y^n. Let $\overline{\mathfrak{b}}_{ij}$ be the i-th block from bottom in the j-th column of Y^n. Let $\mathrm{Color}(\overline{\mathfrak{b}}_{ij})$ be the color of $\overline{\mathfrak{b}}_{ij}$, which is an element in I. Define

$$o(\overline{\mathfrak{b}}_{ij}) := \#\{\overline{\mathfrak{b}}_{rs} \in Y^n |\ \mathrm{Color}(\overline{\mathfrak{b}}_{rs}) = \mathrm{Color}(\overline{\mathfrak{b}}_{ij}),\ (r, s) \prec (i, j)\},$$

where $\prec$ is the lexicographical order; i.e., $(r, s) \prec (i, j)$ if and only if $r < i$ or $(r = i$ and $s < j)$. We define

$$\overline{x}(Y^n) := \sum_{\overline{\mathfrak{b}}_{ij} \in Y^n,\ j > 0} \overline{\mathcal{E}}^{\mathrm{Color}(\overline{\mathfrak{b}}_{i,j})}_{o(\overline{\mathfrak{b}}_{i,j}),\ o(\overline{\mathfrak{b}}_{i,j-1})} \in E_{\overline{\Omega}}(\alpha). \tag{4.3}$$

For $1 \leq i \leq N$, we denote by J_i the subspace of $V(\alpha)$ generated by

$$\{v^{\mathrm{Color}(\overline{\mathfrak{b}}_{ij})}_{o(\overline{\mathfrak{b}}_{ij})} |\ 0 \leq j < l_i\}.$$

By construction, one can show that J_i is invariant under $\overline{x}(Y^n)$ and the representation $(J_i,\ \overline{x}(Y^n)|_{J_i})$ of the quiver $(I, \overline{\Omega})$ is isomorphic to the representation $(V(1-i+k, l_i-i+k),\ \overline{x}(l_i-i+k, 1-i+k))$ of the quiver $(I, \overline{\Omega})$ for $1 \leq i < N$. Here, $\overline{x}(Y^n)|_{J_i}$ is the restriction of $\overline{x}(Y^n)$ on the invariant subspace J_i. Hence $\overline{x}(Y^n)$ is contained in the $G(\alpha)$-orbit $\mathcal{O}_f$ corresponding f, which yields

$$\iota_{\Lambda_k}(X_{Y^n}) = \text{ the closure of the conormal bundle of the } G(\alpha)\text{-orbit of } \overline{x}(Y^n). \tag{4.4}$$

By a direct computation, for $t \in \mathbb{Z}_{\geq 0}$, we have

$$\begin{aligned} \ker(\overline{x}(Y^n))^t &= \bigoplus_{i=1}^{N} \ker(\overline{x}(Y^n)|_{J_i})^t \\ &= \bigoplus_{i=1}^{N} \operatorname{span}_{\mathbb{C}}\{v_{o(\overline{\mathfrak{b}}_{ij})}^{\mathrm{Color}(\overline{\mathfrak{b}}_{ij})} \mid \overline{\mathfrak{b}}_{ij} \in \text{the } i\text{-th row of } Y^n,\ j < t\} \\ &= \operatorname{span}_{\mathbb{C}}\{v_{o(\overline{\mathfrak{b}}_{ij})}^{\mathrm{Color}(\overline{\mathfrak{b}}_{ij})} \mid \overline{\mathfrak{b}}_{ij} \in Y^n,\ j < t\}. \end{aligned} \tag{4.5}$$

In the same manner, we take a Young wall $Y^1 \in \mathcal{Y}^1(\Lambda_k)$ such that $\mathrm{wt}(Y^1) = \alpha$. Denote by X_{Y^1} the image of Y^1 under the crystal isomorphism

$$\mathcal{Y}^1(\Lambda_k) \xrightarrow{\sim} \mathbb{B}(\Lambda_k).$$

Let $\mathfrak{b}_{ij}$ be the i-th block from bottom in the j-th column of Y^1, and $\mathrm{Color}(\mathfrak{b}_{ij})$ the color of $\mathfrak{b}_{ij}$. Set

$$o(\mathfrak{b}_{ij}) := \#\{\mathfrak{b}_{rs} \in Y^1 \mid \mathrm{Color}(\mathfrak{b}_{rs}) = \mathrm{Color}(\mathfrak{b}_{ij}),\ (r,s) \prec (i,j)\},$$

where $\prec$ is the lexicographical order, and define

$$x(Y^1) := \sum_{\mathfrak{b}_{ij} \in Y^1,\ j>0} \mathcal{E}^{\mathrm{Color}(\mathfrak{b}_{i,j-1})}_{o(\mathfrak{b}_{i,j}),\ o(\mathfrak{b}_{i,j-1})} \in E_\Omega(\alpha). \tag{4.6}$$

Then we have

$$\iota_{\Lambda_k}(X_{Y^1}) = \text{ the closure of the conormal bundle of the } G(\alpha)\text{-orbit of } x(Y^1). \tag{4.7}$$

Moreover, we obtain

$$\ker(x(Y^1))^t = \operatorname{span}_{\mathbb{C}}\{v_{o(\mathfrak{b}_{ij})}^{\mathrm{Color}(\mathfrak{b}_{ij})} \mid \mathfrak{b}_{ij} \in Y^1,\ j < t\} \tag{4.8}$$

for $t \in \mathbb{Z}_{\geq 0}$.

Example 4.4. We use the same notations as in Example 3.3. Set

$$\alpha := \Lambda_0 - \mathrm{wt}(b) = 4\alpha_0 + 4\alpha_1 + 4\alpha_2 + 3\alpha_3,$$

and let X be the irreducible component in $\mathbb{B}(\Lambda_0)$ corresponding to b via the crystal isomorphism given in Theorem 4.3. Then we have

$$\begin{aligned} x(Y^1) &= (\mathcal{E}^0_{00} + \mathcal{E}^0_{11} + \mathcal{E}^0_{22}) + (\mathcal{E}^1_{10} + \mathcal{E}^1_{21} + \mathcal{E}^1_{32}) + (\mathcal{E}^2_{00} + \mathcal{E}^2_{11} + \mathcal{E}^2_{22}) + (\mathcal{E}^3_{00} + \mathcal{E}^3_{11} + \mathcal{E}^3_{22}), \\ \overline{x}(Y^n) &= (\overline{\mathcal{E}}^0_{10} + \overline{\mathcal{E}}^0_{21} + \overline{\mathcal{E}}^0_{32}) + (\overline{\mathcal{E}}^1_{00} + \overline{\mathcal{E}}^1_{12} + \overline{\mathcal{E}}^1_{23}) + (\overline{\mathcal{E}}^2_{00} + \overline{\mathcal{E}}^2_{11} + \overline{\mathcal{E}}^2_{33}) + (\overline{\mathcal{E}}^3_{00} + \overline{\mathcal{E}}^3_{22}), \end{aligned}$$

and $\iota_{\Lambda_0}(X)$ is the closure of the conormal bundle of the $G(\alpha)$-orbit of $x(Y^1)$ (resp. $\overline{x}(Y^n)$). However, we note that

$$x(Y^1) + \overline{x}(Y^n) \notin E(\alpha)$$

since $[x(Y^1), \overline{x}(Y^n)] \neq 0$.

5. QUIVER VARIETIES AND THE PERFECT CRYSTALS B^1, B^n

In this section, we give an explanation of the 1-1 correspondence between the geometric realization $\mathbb{B}(\Lambda_k)$ and the path realization of the crystal $B(\Lambda_k)$ associated with the perfect crystals B^1 and B^n, and give a geometric interpretation of the fundamental theorem of perfect crystals in the case of the perfect crystals B^1 and B^n. Let $\alpha \in Q^+$ and let λ be a dominant integral weight of level 1. Choose an irreducible component X in $\mathrm{Irr}\Lambda(\lambda, \alpha)$. For a generic point $\chi = x + \overline{x} \in \iota_\lambda(X)$, we will give an explicit description of the λ-path in B^1 (resp. B^n) corresponding to X using the dimensions of the spaces $\ker x^{i+1}/\ker x^i$ (resp. $\ker \overline{x}^{i+1}/\ker \overline{x}^i$) for $i \geq 0$. For this purpose, we need a couple of lemmas.

Lemma 5.1. *Let X_0 be an irreducible component in $\mathrm{Irr}\Lambda(\alpha)$. Then, for any $\chi = x + \overline{x} \in X_0$ and $k \in \mathbb{Z}_{\geq 0}$, we have*

(a) $\ker(x\overline{x})^k = \ker(\overline{x}x)^k$,

(b) $\ker x^k$ *and* $\ker \overline{x}^k$ *are χ-stable,*

(c) $\ker(x\overline{x})^k$ *is χ-stable.*

Proof. Let $\chi = x + \overline{x} \in X_0$ and $k \in \mathbb{Z}_{\geq 0}$. By (4.1) we have $[x, \overline{x}] = 0$, which yields

$$\chi(\ker x^k) \subset \ker x^k, \ \chi(\ker \overline{x}^k) \subset \ker \overline{x}^k \quad \text{and} \quad \chi(\ker(x\overline{x})^k) \subset \ker(x\overline{x})^k.$$

Our assertion follows from the fact that $\ker(x\overline{x})^k$, $\ker x^k$ and $\ker \overline{x}^k$ are I-graded vector spaces. □

Lemma 5.2. *For each $X_0 \in \mathrm{Irr}\Lambda(\alpha)$, there exists an open subset $U \subset X_0$ such that*

$$\ker x^k \cong \ker {x'}^k \quad \textit{and} \quad \ker \overline{x}^k \cong \ker \overline{x}'^k \tag{5.1}$$

for any $\chi = x + \overline{x}$, $\chi' = x' + \overline{x}' \in U$ and $k \in \mathbb{Z}_{\geq 0}$.

Proof. By Theorem 4.1, there is an open subset $U_1 \subset X_0$ such that $\pi_\Omega(U_1)$ is contained in the $G(\alpha)$-orbit of some element in $E_\Omega(\alpha)$. In the same manner, there is an open subset $U_2 \subset X_0$ such that $\pi_{\overline{\Omega}}(U_2)$ is contained in the $G(\alpha)$-orbit of some element in $E_{\overline{\Omega}}(\alpha)$. Set $U = U_1 \cap U_2 \subset X_0$. Then, by construction, for any $\chi = x + \overline{x}$, $\chi' = x' + \overline{x}' \in U$, there exist $g, \overline{g} \in G(\alpha)$ such that

$$x = gx'g^{-1} \quad \text{and} \quad \overline{x} = \overline{g}\overline{x}'\overline{g}^{-1},$$

which yield, for any $k \in \mathbb{Z}_{\geq 0}$,

$$\ker x^k = \ker(gx'g^{-1})^k = g(\ker {x'}^k) \quad \text{and} \quad \ker \overline{x}^k = \ker(\overline{g}\overline{x}'\overline{g}^{-1})^k = \overline{g}(\ker \overline{x}'^k).$$

□

An element $\chi \in X_0$ in the open subset $U \subset X_0$ in Lemma 5.2 will be called a *generic point.* Thanks to Lemma 5.2, we may consider

$$\underline{\dim}(\ker x^k) \ \text{ and } \ \underline{\dim}(\ker x^{k+1}/\ker x^k) \ (\text{resp. } \underline{\dim}(\ker \overline{x}^k) \ \text{ and } \ \underline{\dim}(\ker \overline{x}^{k+1}/\ker \overline{x}^k))$$

for a generic point $\chi = x + \overline{x}$ in an irreducible component $X_0 \in \mathrm{Irr}\Lambda(\alpha)$. Recall the injective map given in (4.2)

$$\iota_{\Lambda_k} : \mathbb{B}(\Lambda_k) \hookrightarrow \mathbb{B}(\infty)$$

for $0 \le k \le n$. Applying Lemma 5.1 and Lemma 5.2 to (4.6) and (4.3), we obtain the following theorem.

Theorem 5.3. (*cf.* [3]) *Let*

$$\mathbf{p}_k^1 : \mathbb{B}(\Lambda_k) \longrightarrow \mathcal{P}^1(\Lambda_k) \quad (\textit{resp.}\ \mathbf{p}_k^n : \mathbb{B}(\Lambda_k) \longrightarrow \mathcal{P}^n(\Lambda_k)\ )$$

be the unique crystal isomorphism given by Theorem 2.2 and Theorem 4.3, and take an irreducible component $X \in \mathbb{B}(\Lambda_k)$. Then, for a generic point $\chi = x + \overline{x} \in \iota_{\Lambda_k}(X)$, we have

(a)

$$\mathbf{p}_k^1(X) = (\dots, \mathbf{b}_{a_i}, \dots, \mathbf{b}_{a_1}, \mathbf{b}_{a_0}),$$

where $a_i \equiv \dim(\ker x^{i+1} / \ker x^i) - i + k \ (\textit{mod}\ n+1)$ *for all* $i \ge 0$*,*

(b)

$$\mathbf{p}_k^n(X) = (\dots, \overline{\mathbf{b}}_{b_i}, \dots, \overline{\mathbf{b}}_{b_1}, \overline{\mathbf{b}}_{b_0}),$$

where $b_i \equiv 1 - \dim(\ker \overline{x}^{i+1} / \ker \overline{x}^i) + i + k \ (\textit{mod}\ n+1)$ *for all* $i \ge 0$*.*

Proof. Let U be an open subset of $\iota_{\Lambda_k}(X)$ as in Lemma 5.2. By Lemma 5.2, it suffices to show that (a) and (b) hold for some $\chi = x + \overline{x} \in U$.

Let Y^1 be the Young wall in $\mathcal{Y}^1(\Lambda_k)$ corresponding to X under the crystal isomorphism $\mathcal{Y}^1(\Lambda_k) \cong \mathbb{B}(\Lambda_k)$, and $\mathfrak{b}_{ij}$ the i-th block from bottom in the j-th column of Y^1. By (4.7), there exists $\chi = x + \overline{x} \in U$ such that

$$x = g x(Y^1) g^{-1}$$

for some $g \in G(\alpha)$. Then, by the equation (4.8), for $t \in \mathbb{Z}_{\ge 0}$, we have

$$\ker x^t = \ker(g x(Y^1) g^{-1})^t = g(\ker(x(Y^1)^t)) = g\left(\operatorname{span}_{\mathbb{C}}\{v_{\mathfrak{o}(\mathfrak{b}_{ij})}^{\mathrm{Color}(\mathfrak{b}_{ij})} \mid \mathfrak{b}_{ij} \in Y^1,\ j < t\}\right).$$

Let y_t be the t-th column of Y^1 for $t \in \mathbb{Z}_{\ge 0}$. Note that $y_t = \{\mathfrak{b}_{it} \in Y^1 \mid i \ge 1\}$. Then, we have

$$\begin{aligned}
\mathrm{wt}(y_t) &= \sum_{\mathfrak{b}_{it} \in y_t} \alpha_{\mathrm{Color}(\mathfrak{b}_{it})} \\
&= \underline{\dim}\left(\operatorname{span}_{\mathbb{C}}\{v_{o(\mathfrak{b}_{it})}^{\mathrm{Color}(\mathfrak{b}_{it})} \mid \mathfrak{b}_{it} \in y_t\}\right) \\
&= \underline{\dim}\left(\operatorname{span}_{\mathbb{C}}\{v_{o(\mathfrak{b}_{ij})}^{\mathrm{Color}(\mathfrak{b}_{ij})} \mid \mathfrak{b}_{ij} \in Y^1,\ j < t+1\}\right) - \underline{\dim}\left(\operatorname{span}_{\mathbb{C}}\{v_{o(\mathfrak{b}_{ij})}^{\mathrm{Color}(\mathfrak{b}_{ij})} \mid \mathfrak{b}_{ij} \in Y^1,\ j < t\}\right) \\
&= \underline{\dim}(\ker(x(Y^1))^{t+1}) - \underline{\dim}(\ker(x(Y^1))^t) \\
&= \underline{\dim}(\ker x^{t+1}) - \underline{\dim}(\ker x^t) \\
&= \underline{\dim}(\ker x^{t+1} / \ker x^t), \qquad (5.2)
\end{aligned}$$

which implies that the height of y_t is

$$\mathrm{ht}(\mathrm{wt}(y_t)) = \dim(\ker x^{t+1} / \ker x^t).$$

Consequently, the assertion (a) follows from (3.1) and (3.2).

The remaining assertion (b) can be proved in the same manner. $\square$

Combining the crystal isomorphisms (3.1) and (3.3):

$$\mathbf{Y}_k^1 : \mathcal{P}^1(\Lambda_k) \longrightarrow \mathcal{Y}^1(\Lambda_k) \quad \text{and} \quad \mathbf{Y}_k^n : \mathcal{P}^n(\Lambda_k) \longrightarrow \mathcal{Y}^n(\Lambda_k)$$

with (4.7) and (4.4), we have the following proposition, which, together with Theorem 5.3, yields an explicit 1-1 correspondence between $\mathbb{B}(\Lambda_k)$ and $\mathcal{P}^1(\Lambda_k)$ (resp. $\mathcal{P}^n(\Lambda_k)$).

Proposition 5.4. *Let*

$$\mathbf{q}_k^1 : \mathcal{P}^1(\Lambda_k) \longrightarrow \mathbb{B}(\Lambda_k) \;\; (\textit{resp. } \mathbf{q}_k^n : \mathcal{P}^n(\Lambda_k) \longrightarrow \mathbb{B}(\Lambda_k))$$

be the unique crystal isomorphism given by Theorem 2.2 and Theorem 4.3, and take a Λ_k*-path* $\mathbf{p}^1 \in \mathcal{P}^1(\Lambda_k)$ *(resp.* $\mathbf{p}^n \in \mathcal{P}^n(\Lambda_k)$*). Let*

$$\alpha = \Lambda_k - \mathrm{wt}(\mathbf{p}^1) \;\; \textit{and} \;\; X_1 = \mathbf{q}_k^1(\mathbf{p}^1) \quad (\textit{resp. } \beta = \Lambda_k - \mathrm{wt}(\mathbf{p}^n) \;\; \textit{and} \;\; X_n = \mathbf{q}_k^n(\mathbf{p}^n)).$$

Then

(a) $\iota_{\Lambda_k}(X_1)$ *is the closure of the conormal bundle of the* $G(\alpha)$*-orbit of* $x(\mathbf{Y}_k^1(\mathbf{p}^1))$*;*
(b) $\iota_{\Lambda_k}(X_n)$ *is the closure of the conormal bundle of the* $G(\beta)$*-orbit of* $\overline{x}(\mathbf{Y}_k^n(\mathbf{p}^n))$*.*

Recall the fundamental isomorphism theorem of perfect crystals (2.1). From Theorem 4.3, we have the following crystal isomorphisms:

$$\psi_k^1 : \mathbb{B}(\Lambda_k) \xrightarrow{\sim} \mathbb{B}(\Lambda_{k-1}) \otimes B^1,$$
$$\psi_k^n : \mathbb{B}(\Lambda_k) \xrightarrow{\sim} \mathbb{B}(\Lambda_{k+1}) \otimes B^n$$

for $0 \le k \le n$. We would like to give a geometric interpretation to the crystal isomorphisms $\psi_i^1, \; \psi_i^n$ in terms of quiver varieties. To do that, we need a couple of lemmas.

Let V be an I-graded vector space and χ an element of $\mathrm{Hom}(V,V)$. If W is a χ-invariant I-graded subspace of V, then χ can be viewed as an element in $\mathrm{Hom}(V/W, V/W)$ (resp. $\mathrm{Hom}(W,W)$), which is denoted by $\chi|_{V/W}$ (resp. $\chi|_W$).

Lemma 5.5. *Let* $x \in \bigoplus_{i\in I} \mathrm{Hom}(V_{i-1}, V_i)$ *for an* I*-graded vector space* $V := \bigoplus_{i\in I} V_i$*, and set*

$$W := \ker x \;\; \textit{and} \;\; y := x|_{V/W}.$$

Take an element

$$\overline{y} \in \bigoplus_{i\in I} \mathrm{Hom}(V_i/W_i, V_{i-1}/W_{i-1}) \quad \textit{with} \quad [y, \overline{y}] = 0,$$

where W_i *is the* i*-subspace of* W *for* $i \in I$*. Then there exists an element*

$$\overline{x} \in \bigoplus_{i\in I} \mathrm{Hom}(V_i, V_{i-1})$$

such that

$$[x, \overline{x}] = 0 \quad \textit{and} \quad \overline{x}|_{V/W} = \overline{y}.$$

Proof. Let $r = \dim V - \dim W$ and $s = \dim W$. Take an ordered basis for W and extend it to be an ordered basis for V so that the matrix representations of x, y and $\overline{y}$ are given as follows:

$$x = \begin{pmatrix} 0 & A \\ 0 & B \end{pmatrix}, \quad y = B \quad \text{and} \quad \overline{y} = C$$

for some $r \times r$ matrices B and C, and $s \times r$ matrix A. Note that $[B, C] = 0$. Since the matrix

$$\begin{pmatrix} A \\ B \end{pmatrix}$$

has full rank, the following equation

$$\begin{pmatrix} X & Y \\ 0 & 0 \end{pmatrix} \cdot \begin{pmatrix} 0 & A \\ 0 & B \end{pmatrix} = \begin{pmatrix} 0 & AC \\ 0 & 0 \end{pmatrix} \tag{5.3}$$

has a solution. Since $x(V_i) \subset V_{i+1}$ and $\left(\begin{smallmatrix} 0 & AC \\ 0 & 0 \end{smallmatrix}\right)$ maps V_i to V_i, we can choose an $s \times s$ matrix X and an $s \times r$ matrix Y such that $\left(\begin{smallmatrix} X & Y \\ 0 & 0 \end{smallmatrix}\right)$ is a solution of the equation (5.3) and maps V_i to V_{i-1} for $i \in I$. Let

$$\overline{x} = \begin{pmatrix} X & Y \\ 0 & C \end{pmatrix}.$$

By construction, we have

$$\overline{x}(V_i) \subset V_{i-1} \ (i \in I), \qquad \overline{x}|_{V/W} = \overline{y},$$

and

$$[x, \overline{x}] = x\overline{x} - \overline{x}x = \begin{pmatrix} 0 & AC \\ 0 & BC \end{pmatrix} - \begin{pmatrix} 0 & XA + YB \\ 0 & CB \end{pmatrix} = 0.$$

□

Lemma 5.6. *Let U be an open subset of $X_0 \in \mathrm{Irr}\Lambda(\alpha)$ as in Lemma 5.2. Set*

$$\beta = \underline{\dim}(\ker x) \ (\textit{resp. } \gamma = \underline{\dim}(\ker \overline{x}))$$

for $\chi = x + \overline{x} \in U$.

(a) *There exists an irreducible component $X_0' \in \mathrm{Irr}\Lambda(\alpha - \beta)$ such that, for $\chi = x + \overline{x} \in U$,*

$$\phi \circ (\chi|_{V(\alpha)/\ker x}) \circ \phi^{-1} \in X_0', \tag{5.4}$$

where $\phi : V(\alpha)/\ker x \to V(\alpha - \beta)$ is an I-graded vector space isomorphism.

(b) *There exists an irreducible component $X_0'' \in \mathrm{Irr}\Lambda(\alpha - \gamma)$ such that, for $\chi = x + \overline{x} \in U$,*

$$\phi \circ (\chi|_{V(\alpha)/\ker \overline{x}}) \circ \phi^{-1} \in X_0'', \tag{5.5}$$

where $\phi : V(\alpha)/\ker \overline{x} \to V(\alpha - \gamma)$ is an I-graded vector space isomorphism.

Proof. Note that β and γ are well-defined by Lemma 5.2. We first deal with the case (a). For an element $\chi = x + \overline{x} \in U$, let

$$\chi_\phi := \phi \circ (\chi|_{V(\alpha)/\ker x}) \circ \phi^{-1} \in \mathrm{End}(V(\alpha - \beta)),$$

where $\phi : V(\alpha)/\ker x \to V(\alpha - \beta)$ is an I-graded vector space isomorphism. Since $\chi \in \Lambda(\alpha)$, we have

$$\chi_\phi \in \Lambda(\alpha - \beta).$$

Take two elements $\chi = x + \overline{x},\ \chi' = x' + \overline{x}' \in U$, and choose two I-graded vector space isomorphisms $\phi : V(\alpha)/\ker x \to V(\alpha - \beta)$ and $\phi' : V(\alpha)/\ker x' \to V(\alpha - \beta)$. From the properties of U described in the proof of Lemma 5.2, we have

$$x = gx'g^{-1}$$

for some $g \in G(\alpha)$, which yields that $\pi_\Omega(\chi_\phi)$ and $\pi_\Omega(\chi'_{\phi'})$ are in the same $G(\alpha - \beta)$-orbit. Therefore, there exists an irreducible component $X'_0 \in \mathrm{Irr}\Lambda(\alpha - \beta)$ such that

$$\chi_\phi,\ {\chi'}_{\phi'} \in X'_0.$$

Since χ, χ' are arbitrary, our assertion follows.

The remaining case (b) can be proved in the same manner. $\square$

Theorem 5.7. *Let* $X_0 = \iota_{\Lambda_k}(X)$ *for an irreducible component* $X \in \mathrm{Irr}\Lambda(\Lambda_k, \alpha)$*. Set*

$$d = \dim(\ker x) \quad \text{and} \quad \beta = \underline{\dim}(\ker x) \ (\text{resp. } d' = \dim(\ker \overline{x}) \quad \text{and} \quad \gamma = \underline{\dim}(\ker \overline{x}))$$

for a generic point $\chi = x + \overline{x} \in X_0$*.*

(a) *There exists a unique irreducible component* $X' \in \mathrm{Irr}\Lambda(\Lambda_{k-1}, \alpha - \beta)$ *satisfying the following conditions:*
   (i) *there is an open subset* $U \subset X_0$ *such that, for* $\chi = x + \overline{x} \in U$*,*

$$\phi \circ (\chi|_{V(\alpha)/\ker x}) \circ \phi^{-1} \in \iota_{\Lambda_{k-1}}(X'),$$

   where $\phi : V(\alpha)/\ker x \to V(\alpha - \beta)$ *is an* I*-graded vector space isomorphism,*
   (ii) *there is an open subset* $U' \subset \iota_{\Lambda_{k-1}}(X')$ *such that any element* $\chi' \in U'$ *can be written as*

$$\chi' = \phi \circ (\chi|_{V(\alpha)/\ker x}) \circ \phi^{-1},$$

   for some $\chi = x + \overline{x} \in X_0$ *and some* I*-graded vector space isomorphism* $\phi : V(\alpha)/\ker x \to V(\alpha - \beta)$*,*
   (iii) *moreover, we have*

$$\psi_k^1(X) = X' \otimes \mathbf{b}_a \quad \text{and} \quad \mathrm{wt}(\mathbf{b}_a) = \Lambda_k - \Lambda_{k-1} - \mathrm{cl}(\beta),$$

   where $a \equiv d + k$ *(mod* $n + 1$*).*

(b) *There exists a unique irreducible component* $X'' \in \mathrm{Irr}\Lambda(\Lambda_{k+1}, \alpha - \gamma)$ *satisfying the following conditions:*

(i) *there is an open subset* $U \subset X_0$ *such that, for* $\chi = x + \overline{x} \in U$,

$$\phi \circ (\chi|_{V(\alpha)/\ker \overline{x}}) \circ \phi^{-1} \in \iota_{\Lambda_{k+1}}(X''),$$

where $\phi : V(\alpha)/\ker \overline{x} \to V(\alpha - \gamma)$ *is an* I*-graded vector space isomorphism,*

(ii) *there is an open subset* $U'' \subset \iota_{\Lambda_{k+1}}(X'')$ *such that any element* $\chi'' \in U''$ *can be written as*

$$\chi'' = \phi \circ (\chi|_{V(\alpha)/\ker \overline{x}}) \circ \phi^{-1},$$

for some $\chi = x + \overline{x} \in X_0$ *and some* I*-graded vector space isomorphism* $\phi : V(\alpha)/\ker \overline{x} \to V(\alpha - \gamma)$,

(iii) *moreover, we have*

$$\psi_k^n(X) = X'' \otimes \overline{\mathbf{b}}_b \quad \text{and} \quad \mathrm{wt}(\overline{\mathbf{b}}_b) = \Lambda_k - \Lambda_{k+1} - \mathrm{cl}(\gamma),$$

where $b \equiv 1 - d' + k \ (mod\ n+1)$.

Proof. We first deal with the case (a) of the crystal isomorphism $\psi_k^1 : \mathbb{B}(\Lambda_k) \to \mathbb{B}(\Lambda_{k-1}) \otimes B^1$. Let Y be the Young wall in $\mathcal{Y}^1(\Lambda_k)$ corresponding to X and Y' the Young wall obtained by removing the 0-th column from Y. Then Y' can be viewed as an element in $\mathcal{Y}^1(\Lambda_{k-1})$. Take the irreducible component X' in $\mathbb{B}(\Lambda_{k-1})$ corresponding to Y'. By Theorem 5.3 and (5.2), we have

$$X' \in \mathrm{Irr}\Lambda(\Lambda_{k-1}, \alpha - \beta) \quad \text{and} \quad \psi_k^1(X) = X' \otimes \mathbf{b}_a,$$

where $a \equiv d + k \pmod{n+1}$ and $\mathrm{wt}(\mathbf{b}_a) = \Lambda_k - \Lambda_{k-1} - \mathrm{cl}(\beta)$.

Let U be an open subset of X_0 as in Lemma 5.6, and take an element $\chi = x + \overline{x} \in U$. Since $x(Y)|_{V(\alpha)/\ker(x(Y))}$ is naturally identified with $x(Y')$ and x is contained in the $G(\alpha)$-orbit of $x(Y)$, by Lemma 5.6, we have

$$\phi \circ (\chi|_{V(\alpha)/\ker x}) \circ \phi^{-1} \in \iota_{\Lambda_{k-1}}(X')$$

for an I-graded vector space isomorphism $\phi : V(\alpha)/\ker x \to V(\alpha - \beta)$.

Take an element $\chi' = x' + \overline{x}'$ in an open subset U' of $\iota_{\Lambda_{k-1}}(X')$ given as in Lemma 5.2. Then x' can be written as

$$x' = g x(Y') g^{-1}$$

for some $g \in G(\alpha - \beta)$, which yields that there is x in the $G(\alpha)$-orbit of $x(Y)$ such that

$$\phi \circ (x|_{V(\alpha)/\ker x}) \circ \phi^{-1} = x'$$

for some I-graded vector space isomorphism $\phi : V(\alpha)/\ker x \to V(\alpha - \beta)$. The assertion (ii) follows from Lemma 5.5.

The remaining case (b), $\psi_k^n : \mathbb{B}(\Lambda_k) \to \mathbb{B}(\Lambda_{k+1}) \otimes B^n$, can be proved in the same manner. □

Example 5.8. We use the same notations as in Example 4.4. Let $X_0 = \iota_{\Lambda_0}(X) \in \mathrm{Irr}\Lambda(\alpha)$. By Theorem 4.1, it suffices to consider a generic point in the fiber $\pi_{\overline{\Omega}}^{-1}(\overline{x}(Y^n)) \subset X_0$. By [14, Section 12.8, Proposition 15.5], we have

$$\begin{aligned}
\pi_{\overline{\Omega}}^{-1}(\overline{x}(Y^n)) &= \{\chi \in X_0 |\ \pi_{\overline{\Omega}}(\chi) = \overline{x}(Y^n)\} \\
&= \{x + \overline{x}(Y^n) |\ x \in E_\Omega(\alpha),\ [x, \overline{x}(Y^n)] = 0\} \\
&= \{x + \overline{x}(Y^n) \mid x = x(a_1, \ldots, a_{13}),\ a_1, \ldots, a_{13} \in \mathbb{C}\}.
\end{aligned}$$

Here,

$$x(a_1, \ldots, a_{13}) = \begin{pmatrix} 0 & 0 & 0 & x_0 \\ x_1 & 0 & 0 & 0 \\ 0 & x_2 & 0 & 0 \\ 0 & 0 & x_3 & 0 \end{pmatrix} \in E_\Omega(\alpha),$$

$$x_0 = \begin{pmatrix} a_1 & a_2 & a_3 \\ 0 & 0 & 0 \\ a_4 & 0 & a_5 \\ a_6 & 0 & 0 \end{pmatrix}, \qquad x_1 = \begin{pmatrix} 0 & a_1 & a_2 & a_3 \\ 0 & a_4 & 0 & a_5 \\ 0 & a_6 & 0 & 0 \\ 0 & a_{10} & 0 & a_7 \end{pmatrix},$$

$$x_2 = \begin{pmatrix} 0 & a_2 & a_3 & 0 \\ 0 & 0 & a_5 & 0 \\ a_6 & a_8 & a_{11} & a_9 \\ 0 & 0 & a_7 & 0 \end{pmatrix}, \qquad x_3 = \begin{pmatrix} 0 & a_2 & 0 & 0 \\ a_4 & a_{12} & a_5 & a_{13} \\ a_6 & a_8 & 0 & a_9 \end{pmatrix}$$

for $a_1, \ldots, a_{13} \in \mathbb{C}$. Let

$$x := x(a_1, \ldots, a_{13}), \qquad \overline{x} := \overline{x}(Y^n),$$

and consider $a_1, \ldots, a_{13}$ as indeterminates. Then we have

$$\dim(\ker x^k) = \begin{cases} 0 & \text{if } k = 0, \\ 2 + k & \text{if } 1 \le k \le 12, \\ 15 & \text{otherwise,} \end{cases} \quad \text{and} \quad \dim(\ker \overline{x}^k) = \begin{cases} 0 & \text{if } k = 0, \\ 4 & \text{if } k = 1, \\ 8 & \text{if } k = 2, \\ 11 & \text{if } k = 3, \\ 14 & \text{if } k = 4, \\ 15 & \text{otherwise.} \end{cases}$$

Hence we obtain

$$\begin{aligned}
&\mathbf{p}_0^1(X) = (\ldots, \mathbf{b}_3, \mathbf{b}_1, \mathbf{b}_2, \mathbf{b}_3, \mathbf{b}_0, \mathbf{b}_1, \mathbf{b}_2, \mathbf{b}_3, \mathbf{b}_0, \mathbf{b}_1, \mathbf{b}_2, \mathbf{b}_3, \mathbf{b}_0, \mathbf{b}_3), \\
&\mathbf{p}_0^n(X) = (\ldots, \overline{\mathbf{b}}_3, \overline{\mathbf{b}}_2, \overline{\mathbf{b}}_0, \overline{\mathbf{b}}_1, \overline{\mathbf{b}}_0, \overline{\mathbf{b}}_2, \overline{\mathbf{b}}_1).
\end{aligned}$$

6. Quiver Varieties and Adjoint Crystals

In this section, we will prove the main theorem of this paper, Theorem 6.3, which shows that there exists an explicit crystal isomorphism between the geometric realization $\mathbb{B}(\Lambda_0)$ and the path realization $\mathcal{P}^{\mathrm{ad}}(\Lambda_0)$ of $B(\Lambda_0)$ arising from the adjoint crystal B^{ad}. By Theorem 2.2 and Theorem 4.3, we have the crystal isomorphism

$$\mathbf{p}^{\mathrm{ad}} : \mathbb{B}(\Lambda_0) \longrightarrow \mathcal{P}^{\mathrm{ad}}(\Lambda_0).$$

Let $\alpha \in Q^+$ and let X be an irreducible component of $\mathrm{Irr}\Lambda(\Lambda_0, \alpha)$. For a generic point $\chi = x + \overline{x} \in \iota_{\Lambda_0}(X)$, we will give an explicit description of the Λ_0-path $\mathbf{p}^{\mathrm{ad}}(X)$ in terms of dimension vectors of $\ker(x\overline{x})^{k+1}/\ker(x\overline{x})^k$ for $k \geq 0$.

Let $\alpha, \beta \in Q^+$ with $\beta \leq \alpha$. Consider the diagram given in [14]:

$$\Lambda(\alpha-\beta) \xleftarrow{\pi} \Lambda(\alpha-\beta) \times \Lambda(\beta) \xleftarrow{p_1} F' \xrightarrow{p_2} F'' \xrightarrow{p_3} \Lambda(\alpha), \tag{6.1}$$

where F'' is the variety of all pairs (χ, W) such that

(a) $\chi \in \Lambda(\alpha)$,
(b) W is a χ-stable subspace of $V(\alpha)$ with $\underline{\dim} W = \beta$,

and F' is the variety of all quadruples (χ, W, f, g) such that

(a) $(\chi, W) \in F''$,
(b) $f = (f_i)_{i\in I}, g = (g_i)_{i \in I}$ give an exact sequence

$$0 \longrightarrow V_i(\beta) \xrightarrow{f_i} V_i(\alpha) \xrightarrow{g_i} V_i(\alpha-\beta) \longrightarrow 0 \quad (i \in I)$$

such that $\mathrm{im} f = W$.

Then we have

$$p_1(\chi, W, f, g) = (\tilde{g} \circ (\chi|_{V(\alpha)/W}) \circ \tilde{g}^{-1},\ f^{-1} \circ (\chi|_W) \circ f),$$

where $\tilde{g} : V(\alpha)/W \to V(\alpha-\beta)$ is the I-graded vector space isomorphism induced by g,

$$p_2(\chi, W, f, g) = (\chi, W), \quad p_3(\chi, W) = \chi,$$

and π is the natural first projection. Note that p_2 is a $G(\alpha-\beta) \times G(\beta)$-principal bundle and an open map.

Let U be an open subset of $X_0 \in \mathrm{Irr}\Lambda(\alpha)$ as in Lemma 5.2, and $\beta = \underline{\dim}(\ker x)$ for $\chi = x + \overline{x} \in U$. Define the map $\imath : U \to F''$ by

$$\imath(\chi) = (\chi,\ \ker x)$$

for $\chi = x + \overline{x} \in U$. Note that $p_3 \circ \imath = \mathrm{id}|_U$. By Lemma 5.6, there exists an irreducible component $X_0' \in \mathrm{Irr}\Lambda(\alpha - \beta)$ such that, for any $\chi = x + \overline{x} \in U$,

$$\phi \circ (\chi|_{V(\alpha)/\ker x}) \circ \phi^{-1} \in X_0',$$

where $\phi : V(\alpha)/\ker x \to V(\alpha-\beta)$ is an I-graded vector space isomorphism. Given an open subset $U' \subset \Lambda(\alpha-\beta)$ with $U' \cap X_0' \neq \emptyset$, by Lemma 5.5,

$$\tilde{U} := \imath^{-1} \circ p_2 \circ p_1^{-1} \circ \pi^{-1}(U')$$

is a nonempty open subset of X_0. Therefore, given an open subset $U' \subset X'_0$, there exists an open subset $\tilde{U} \subset X_0$ such that, for any element $\chi = x + \overline{x} \in \tilde{U}$,

$$\phi \circ (\chi|_{V(\alpha)/\ker x}) \circ \phi^{-1} \in U'$$

for some I-graded vector space isomorphism $\phi : V(\alpha)/\ker x \to V(\alpha - \beta)$.

In the same manner, let $\gamma = \underline{\dim}(\ker \overline{x})$, and consider the diagram

$$\Lambda(\alpha-\gamma) \xleftarrow{\pi} \Lambda(\alpha-\gamma) \times \Lambda(\gamma) \xleftarrow{p_1} F' \xrightarrow{p_2} F'' \xrightarrow{p_3} \Lambda(\alpha).$$

Define the map $\overline{\imath} : U \to F''$ by

$$\overline{\imath}(\chi) = (\chi, \ker \overline{x})$$

for $\chi = x + \overline{x} \in U$, and let X''_0 be an irreducible component as in Lemma 5.6. Then one can deduce that, given an open subset $U' \subset X''_0$, there exists an open subset $\tilde{U} \subset X_0$ such that, for any element $\chi = x + \overline{x} \in \tilde{U}$,

$$\phi \circ (\chi|_{V(\alpha)/\ker \overline{x}}) \circ \phi^{-1} \in U'$$

for some I-graded vector space isomorphism $\phi : V(\alpha)/\ker \overline{x} \to V(\alpha - \beta)$. Consequently, we have the following lemma.

Lemma 6.1. *With the same notations as in Lemma 5.6, we have the following.*

(a) *Given an open subset $U' \subset X'_0$, there exists an open subset $\tilde{U} \subset X_0$ such that, for any element $\chi = x + \overline{x} \in \tilde{U}$,*

$$\phi \circ (\chi|_{V(\alpha)/\ker x}) \circ \phi^{-1} \in U' \tag{6.2}$$

for some I-graded vector space isomorphism $\phi : V(\alpha)/\ker x \to V(\alpha - \beta)$.

(b) *Given an open subset $U'' \subset X''_0$, there exists an open subset $\tilde{U} \subset X_0$ such that, for any element $\chi = x + \overline{x} \in \tilde{U}$,*

$$\phi \circ (\chi|_{V(\alpha)/\ker \overline{x}}) \circ \phi^{-1} \in U'' \tag{6.3}$$

for some I-graded vector space isomorphism $\phi : V(\alpha)/\ker \overline{x} \to V(\alpha - \gamma)$.

Combining Lemma 6.1 with Lemma 5.6, we have the following lemma.

Lemma 6.2. *Let $\alpha \in Q^+$. For each $X_0 \in \mathrm{Irr}\Lambda(\alpha)$, there exists an open subset $U \subset X_0$ such that*

$$\ker(x\overline{x})^k \cong \ker(x'\overline{x}')^k \quad \text{and} \quad \ker x(x\overline{x})^k \cong \ker x'(x'\overline{x}')^k \tag{6.4}$$

for any $\chi = x + \overline{x}, \chi' = x' + \overline{x}' \in U$ and $k \in \mathbb{Z}_{\geq 0}$.

Proof. Since the case that $\mathrm{ht}(\alpha) = 0$ is trivial, we may assume $\mathrm{ht}(\alpha) > 0$. Let U_0 be an open subset of X_0 as in Lemma 5.2, and $\beta = \underline{\dim}(\ker x)$ for $\chi = x + \overline{x} \in U_0$. Take the irreducible component $X'_0 \in \mathrm{Irr}\Lambda(\alpha - \beta)$ given in Lemma 5.6, and choose an open subset U'_0 of X'_0 satisfying the conditions of Lemma 5.2. Let $\gamma = \underline{\dim}(\ker \overline{x})$ for $\chi = x + \overline{x} \in U'_0$. By Lemma 6.1, there exists an open subset $\tilde{U}_0 \subset X_0$ such that, for any element $\chi = x + \overline{x} \in \tilde{U}_0$,

$$\phi \circ (\chi|_{V(\alpha)/\ker x}) \circ \phi^{-1} \in U'_0$$

for some I-graded vector space isomorphism $\phi : V(\alpha)/\ker x \to V(\alpha-\beta)$. Set $\hat{U}_0 = U_0 \cap \tilde{U}_0$. Then, for any $\chi = x + \overline{x} \in \hat{U}_0$, we have

$$\begin{aligned}\underline{\dim}\ker x\overline{x} &= \underline{\dim}(\ker x) + \underline{\dim}(\ker x\overline{x}/\ker x)\\ &= \underline{\dim}(\ker x) + \underline{\dim}\{v \in V(\alpha)/\ker x \mid \overline{x}|_{V(\alpha)/\ker x}(v) = 0\}\\ &= \underline{\dim}(\ker x) + \underline{\dim}(\ker \overline{x}|_{V(\alpha)/\ker x})\\ &= \beta + \gamma.\end{aligned}$$

Let us take the irreducible component $X_0'' \in \mathrm{Irr}\Lambda(\alpha-\beta-\gamma)$ associated with X_0', which is given as in Lemma 5.6. By the induction hypothesis, there exists an open subset $U'' \subset X_0''$ satisfying (6.4). Applying Lemma 6.1 to X_0'', X_0' and X_0, there exists an open subset $\tilde{U} \subset X_0$ such that, for any $\chi = x + \overline{x} \in \hat{U}$,

$$\phi \circ (\chi|_{V(\alpha)/\ker x\overline{x}}) \circ \phi^{-1} \in U'' \tag{6.5}$$

for some I-graded vector space isomorphism $\phi : V(\alpha)/\ker x\overline{x} \to V(\alpha-\beta-\gamma)$. Let $U = \tilde{U} \cap \hat{U}_0$, then, by construction, U holds the condition (6.4). □

An element $\chi \in X_0$ in the open subset $U \subset X_0$ satisfying (5.1) and (6.4) is called a *generic point*. Note that, for $\chi = x + \overline{x}$ in an irreducible component $X_0 \in \mathrm{Irr}\Lambda(\alpha)$, since $[x, \overline{x}] = 0$, we have

$$\dim(\ker(x|_{\ker(x\overline{x})^{k+1}/\ker(x\overline{x})^k})) = \dim(\ker x(x\overline{x})^k) - \dim(\ker(x\overline{x})^k),$$

where $x|_{\ker(x\overline{x})^{k+1}/\ker(x\overline{x})^k}$ is the linear map in $\mathrm{End}(\ker(x\overline{x})^{k+1}/\ker(x\overline{x})^k)$ induced by x. Thanks to Lemma 6.2, one can talk about

$$\underline{\dim}\ker(x\overline{x})^k, \quad \underline{\dim}\ker(x\overline{x})^{k+1}/\ker(x\overline{x})^k \quad \text{and} \quad \dim(\ker(x|_{\ker(x\overline{x})^{k+1}/\ker(x\overline{x})^k}))$$

for a generic point $\chi = x + \overline{x}$ in an irreducible component $X_0 \in \mathrm{Irr}\Lambda(\alpha)$ and $k \in \mathbb{Z}_{\geq 0}$.

Finally, we are ready to state the main theorem in this paper.

Theorem 6.3. *Let*

$$\mathbf{p}^{\mathrm{ad}} : \mathbb{B}(\Lambda_0) \longrightarrow \mathcal{P}^{\mathrm{ad}}(\Lambda_0)$$

be the unique crystal isomorphism given by Theorem 2.2 and Theorem 4.3, and take an irreducible component $X \in \mathbb{B}(\Lambda_0)$. For a generic point $\chi = x + \overline{x} \in \iota_{\Lambda_0}(X)$ and $k \in \mathbb{Z}_{\geq 0}$, let

$$\begin{aligned}\theta_k &= \underline{\dim}(\ker(x\overline{x})^{k+1}/\ker(x\overline{x})^k),\\ c_k &\equiv \dim(\ker(x|_{\ker(x\overline{x})^{k+1}/\ker(x\overline{x})^k})) \ (\mathit{mod}\ n+1),\end{aligned}$$

where $0 \leq c_k \leq n$. Then we have

$$\mathbf{p}^{\mathrm{ad}}(X) = (\ldots, p_k, \ldots, p_1, p_0), \tag{6.6}$$

where

$$p_k = \begin{cases} \mathsf{b}_{-\mathrm{cl}(\theta_k)} & \text{if } \mathrm{cl}(\theta_k) \neq 0,\\ h_{c_k} & \text{if } \mathrm{cl}(\theta_k) = 0 \text{ and } c_k \neq 0,\\ \emptyset & \text{otherwise.}\end{cases}$$

Proof. Let $X_0 = \iota_{\Lambda_0}(X)$ with $\mathrm{wt}(X) = \Lambda_0 - \alpha$ for some $\alpha \in Q^+$. We will use induction on $\mathrm{ht}(\alpha)$. Since the case that $\mathrm{ht}(\alpha) = 0$ is trivial, we may assume $\alpha \neq 0$. Note that

$$\begin{aligned}&\theta_k = \underline{\dim}(\ker(x\overline{x})^{k+1}) - \underline{\dim}(\ker(x\overline{x})^k),\\ &c_k \equiv \dim(\ker x(x\overline{x})^k) - \dim(\ker(x\overline{x})^k) \pmod{n+1}\end{aligned}$$

for a generic point $\chi = x + \overline{x} \in X_0$ and $k \in \mathbb{Z}_{\geq 0}$.

Let $\beta = \underline{\dim}(\ker x)$ for a generic point $\chi = x + \overline{x} \in X_0$, and choose the irreducible component $X_0' \in \mathrm{Irr}\Lambda(\alpha - \beta)$ associated with X_0 as in Lemma 5.6. Similarly, let $\gamma = \underline{\dim}(\ker \overline{x})$ for a generic point $\chi = x + \overline{x} \in X_0'$, and take the irreducible component $X_0'' \in \mathrm{Irr}\Lambda(\alpha - \beta - \gamma)$ associated with X_0' as in Lemma 5.6. By Theorem 5.7, we have

$$\psi_0^1(X_0) = X_0' \otimes \mathbf{b}_a \quad \text{and} \quad \psi_n^n(X_0') = X_0'' \otimes \overline{\mathbf{b}}_b$$

for some $\mathbf{b}_a \in B^1$, $\overline{\mathbf{b}}_b \in B^n$. From (2.1) and Theorem 4.2, we have

$$\psi_0^{\mathrm{ad}} : \mathbb{B}(\Lambda_0) \xrightarrow{\sim} \mathbb{B}(\Lambda_0) \otimes B^{\mathrm{ad}}.$$

Then, it follows from the crystal isomorphism (2.2) and Theorem 5.7 that

$$\psi_0^{\mathrm{ad}}(X_0) = X_0'' \otimes \mathfrak{p}^{\mathrm{ad}}(\overline{\mathbf{b}}_b \otimes \mathbf{b}_a).$$

By the induction hypothesis, there is an open subset $U'' \subset X_0''$ satisfying (6.6). By Lemma 6.1 and (6.5), there is an open subset $\tilde{U} \subset X_0$ such that, for any $\chi = x + \overline{x} \in \tilde{U}$,

$$\phi \circ (\chi|_{V(\alpha)/\ker x\overline{x}}) \circ \phi^{-1} \in U''$$

for some isomorphism $\phi : V(\alpha)/\ker x\overline{x} \to V(\alpha - \beta - \gamma)$.

On the other hand, by Lemma 6.2, there exists an open subset $\hat{U} \subset X_0$ satisfying (6.4). Set

$$U = \hat{U} \cap \tilde{U}$$

and choose an element $\chi = x + \overline{x} \in U$. Suppose that $\mathrm{wt}(\overline{\mathbf{b}}_b \otimes \mathbf{b}_a) \neq 0$. Then, by Theorem 5.7 and (2.2), since

$$\begin{aligned}\mathrm{wt}(\overline{\mathbf{b}}_b \otimes \mathbf{b}_a) &= \mathrm{wt}(\overline{\mathbf{b}}_b) + \mathrm{wt}(\mathbf{b}_a)\\ &= \Lambda_n - \Lambda_0 - \mathrm{cl}(\gamma) + \Lambda_0 - \Lambda_n - \mathrm{cl}(\beta)\\ &= -\mathrm{cl}(\beta + \gamma)\\ &= -\mathrm{cl}(\underline{\dim}(\ker x\overline{x}))\\ &= -\mathrm{cl}(\theta_0),\end{aligned}$$

we obtain

$$\mathfrak{p}^{\mathrm{ad}}(\overline{\mathbf{b}}_b \otimes \mathbf{b}_a) = \mathsf{b}_{\mathrm{wt}(\overline{\mathbf{b}}_b \otimes \mathbf{b}_a)} = \mathsf{b}_{-\mathrm{cl}(\theta_0)}.$$

Suppose $\mathrm{wt}(\overline{\mathbf{b}}_b \otimes \mathbf{b}_a) = 0$ and $a \neq n+1$. Then, by Theorem 5.7 and (2.2), we have

$$\mathfrak{p}^{\mathrm{ad}}(\overline{\mathbf{b}}_b \otimes \mathbf{b}_a) = h_a$$

and $a \equiv \dim x \pmod{n+1}$, which implies that $a = c_0$. In the same manner, if $\mathrm{wt}(\overline{\mathbf{b}}_b \otimes \mathbf{b}_a) = 0$ and $a = n+1$, we have

$$\mathfrak{p}^{\mathrm{ad}}(\overline{\mathbf{b}}_b \otimes \mathbf{b}_a) = \emptyset.$$

Since, for an arbitrary isomorphism $\phi : V(\alpha)/\ker x\overline{x} \to V(\alpha - \beta - \gamma)$,

$$\begin{aligned}\underline{\dim}(\ker(x\overline{x})^{k+1}) &= \underline{\dim}(\ker(x\overline{x})^{k+1}/\ker(x\overline{x})) + \underline{\dim}(\ker(x\overline{x}))\\ &= \underline{\dim}(\ker(x\overline{x})|_{V(\alpha)/\ker(x\overline{x})})^k + \underline{\dim}(\ker(x\overline{x}))\\ &= \underline{\dim}\ker(\phi \circ (x\overline{x})|_{V(\alpha)/\ker(x\overline{x})} \circ \phi^{-1})^k + \underline{\dim}(\ker(x\overline{x}))\\ &= \underline{\dim}\ker((\phi \circ x|_{V(\alpha)/\ker(x\overline{x})} \circ \phi^{-1})(\phi \circ \overline{x}|_{V(\alpha)/\ker(x\overline{x})} \circ \phi^{-1}))^k + \underline{\dim}(\ker(x\overline{x})),\end{aligned}$$

our assertion follows from a standard induction argument. □

The following corollary, which is an immediate consequence of Theorem 5.7 and Theorem 6.3, can be regarded as a geometric interpretation of the fundamental isomorphism theorem for perfect crystals

$$\psi_0^{\mathrm{ad}} : \mathbb{B}(\Lambda_0) \xrightarrow{\sim} \mathbb{B}(\Lambda_0) \otimes B^{\mathrm{ad}}.$$

Corollary 6.4. *Let $X_0 = \iota_{\Lambda_0}(X)$ for some $X \in \mathrm{Irr}\Lambda(\Lambda_0, \alpha)$. For a generic point $\chi = x + \overline{x} \in X_0$, set*

$$\theta = \underline{\dim}(\ker(x\overline{x})) \quad \textit{and} \quad c = \dim(\ker x).$$

Then there exists a unique irreducible component $X' \in \mathrm{Irr}\Lambda(\Lambda_0, \alpha - \theta)$ satisfying the following conditions:

(a) *there is an open subset $U \subset X_0$ such that, for $\chi = x + \overline{x} \in U$,*

$$\phi \circ (\chi|_{V(\alpha)/\ker x\overline{x}}) \circ \phi^{-1} \in \iota_{\Lambda_0}(X'),$$

where $\phi : V(\alpha - \theta) \to V(\alpha)/\ker x\overline{x}$ is an I-graded vector space isomorphism,

(b) *there is an open subset $U' \subset \iota_{\Lambda_0}(X')$ such that any element $\chi' \in U'$ can be written as*

$$\chi' = \phi \circ (\chi|_{V(\alpha)/\ker x\overline{x}}) \circ \phi^{-1}$$

for some $\chi = x + \overline{x} \in X_0$ and some I-graded vector space isomorphism $\phi : V(\alpha)/\ker x\overline{x} \to V(\alpha - \theta)$,

(c) *moreover, we have*

$$\psi_0^{\mathrm{ad}}(X) = X' \otimes p,$$

where

$$p = \begin{cases} \mathsf{b}_{-\mathrm{cl}(\theta)} & \textit{if } \mathrm{cl}(\theta) \neq 0,\\ h_c & \textit{if } \mathrm{cl}(\theta) = 0 \textit{ and } c \neq 0,\\ \emptyset & \textit{otherwise.}\end{cases}$$

Example 6.5. We use the same notations as in Example 5.8. Let $W_i = \ker(x\overline{x})^i$ for $i \in \mathbb{Z}_{\geq 0}$. Then we have

$$\underline{\dim}(W_i) = \begin{cases} 0 & \text{if } i = 0, \\ \alpha_0 + 2\alpha_1 + 2\alpha_2 + \alpha_3 & \text{if } i = 1, \\ 2\alpha_0 + 3\alpha_1 + 3\alpha_2 + 2\alpha_3 & \text{if } i = 2, \\ 3\alpha_0 + 4\alpha_1 + 4\alpha_2 + 3\alpha_3 & \text{if } i = 3, \\ 4\alpha_0 + 4\alpha_1 + 4\alpha_2 + 3\alpha_3 & \text{otherwise,} \end{cases}$$

and

$$\dim(\ker(x|_{W_{i+1}/W_i})) = \dim(\ker x(x\overline{x})^i) - \dim(\ker(x\overline{x})^i) = \begin{cases} 3 & \text{if } i = 0, \\ 1 & \text{if } i = 1, \\ 1 & \text{if } i = 2, \\ 0 & \text{otherwise.} \end{cases}$$

By Theorem 6.3, we have

$$\mathbf{p}^{\mathrm{ad}}(X) = (\ldots, \emptyset, \emptyset, \mathsf{b}_{\alpha_1+\alpha_2+\alpha_3}, h_1, h_1, \mathsf{b}_{-\alpha_1-\alpha_2}).$$

References

[1] G. Benkart, I. Frenkel, S.-J. Kang, and H. Lee, *Level 1 perfect crystals and path realizations of basic representations at* $q = 0$, Int. Math. Res. Not. 2006, Art. ID 10312, 1–28.

[2] E. Date, M. Jimbo, A. Kuniba, T. Miwa, and M. Okado, *Paths, Maya diagrams and representations of* $\widehat{sl}(r, \mathbb{C})$, Adv. Stud. Pure Math. **19** (1989), 149–191.

[3] I. B. Frenkel and A. Savage, *Bases of representations of type A affine Lie algebras via quiver varieties and statistical mechanics*, Int. Math. Res. Not. 2003, no. 28, 1521–1547.

[4] J. Hong, S.-J. Kang, *Introduction to Quantum Groups and Crystal Bases*, Graduate Studies in Mathematics, **42**. American Mathematical Society, Providence, RI, 2002.

[5] M. Jimbo, K. C. Misra, T. Miwa, and M. Okado, *Combinatorics of representations of* $U_q(\hat{sl}(n))$ *at* $q = 0$, Comm. Math. Phys. **136** (1991), no. 3, 543–566.

[6] S.-J. Kang, *Crystal bases for quantum affine algebras and combinatorics of Young walls*, Proc. London Math. Soc. (3) **86** (2003), no. 1, 29–69.

[7] S.-J. Kang, M. Kashiwara, K. C. Misra, T. Miwa, T. Nakashima, and A. Nakayashiki, *Affine Crystals and Vertex Models*, Infinite Analysis, Part A,B(Kyoto, 1991), Adv. Ser. Math. Phys. **16**, World Scientific Publishing Co. Inc., River Edge, NJ, 1992, 449–484.

[8] ———, *Perfect crystals of quantum affine Lie algebras*, Duke Math. J. **68** (1992), no. 3, 499–607.

[9] M. Kashiwara, *Crystalizing the q-analogue of universal enveloping algebras*, Comm. Math. Phys. **133** (1990), no. 2, 249–260.

[10] ———, *On crystal bases of the q-analogue of universal enveloping algebras*, Duke Math. J. **63** (1991), no. 2, 465–516.

[11] ———, *The crystal base and Littelmann's refined Demazure character formula*, Duke Math. J. **71** (1993), no. 3, 839–858.

[12] M. Kashiwara and Y. Saito, *Geometric construction of crystal bases*, Duke Math. J. **89** (1997), no. 1, 9–36.

[13] G. Lusztig, *Canonical bases arising from quantized enveloping algebras*, J. Amer. Math. Soc. **3** (1990), no. 2, 447–498.

[14] ———, *Quivers, perverse sheaves, and quantized enveloping algebras*, J. Amer. Math. Soc. **4** (1991), no. 2, 365–421.

[15] K.C. Misra and T. Miwa, *Crystal base for the basic representation of* $U_q(\hat{sl}(n))$, Comm. Math. Phys. **134** (1990), no. 1, 79–88.

[16] H. Nakajima, *Instantons on ALE spaces, quiver varieties, and Kac-Moody algebras*, Duke Math. J. **76** (1994), no. 2, 365–416.

[17] ———, *Quiver varieties and Kac-Moody algebras*, Duke Math. J. **91** (1998), no. 3, 515–560.

[18] Y. Saito, *Crystal bases and quiver varieties*, Math. Ann. **324** (2002), no. 4, 675–688.

[19] A. Savage, *Geometric and combinatorial realizations of crystal graphs*, Algebr. Represent. Theory **9** (2006), no. 2, 161–199.

Department of Mathematical Sciences and Research Institute of Mathematics, Seoul National University, Gwanak-ro 599, Gwanak-gu, Seoul 151-747, Korea

E-mail address: `sjkang@math.snu.ac.kr`

Department of Mathematical Sciences, Seoul National University, Gwanak-ro 599, Gwanak-gu, Seoul 151-747, Korea

E-mail address: `pwy@snu.ac.kr`